

BEREZIN-TOEPLITZ QUANTIZATION FOR COMPACT KÄHLER MANIFOLDS. A REVIEW OF RESULTS

MARTIN SCHLICHENMAIER

ABSTRACT. This article is a review on Berezin-Toeplitz operator and Berezin-Toeplitz deformation quantization for compact quantizable Kähler manifolds. The basic objects, concepts, and results are given. This concerns the correct semi-classical limit behaviour of the operator quantization, the unique Berezin-Toeplitz deformation quantization (star product), covariant and contravariant Berezin symbols, and Berezin transform. Other related objects and constructions are also discussed.

CONTENTS

1. Introduction	2
2. The set-up of geometric quantization	4
2.1. Kähler manifolds	4
2.2. Poisson algebra	4
2.3. Quantum line bundles	4
2.4. Example: The Riemann sphere	5
2.5. Example: The complex projective space	5
2.6. Example: The torus	5
2.7. Example: The unit disc and Riemann surfaces	6
2.8. Consequences of quantizability	6
2.9. Embedding into projective space	6
3. Berezin-Toeplitz operators	7
3.1. Tensor powers of the quantum line bundle	7
3.2. Approximation results	8
3.3. Further properties	9
3.4. Strict quantization	10
3.5. Relation to geometric quantization	10
3.6. L_α approximation	11
3.7. The noncompact situation	11
4. Berezin-Toeplitz deformation quantization	11
4.1. Definition of star products	12
4.2. Existence of star products	13
4.3. Equivalence and classification of star products	13
4.4. Star products with separation of variables	13

Date: December 30, 2009.

1991 *Mathematics Subject Classification.* Primary: 58F06, 53D55; Secondary: 58G15, 53C55, 32C17, 81S10.

4.5.	Berezin-Toeplitz star product	14
4.6.	Star product of geometric quantization	15
4.7.	Trace for the BT star product	15
4.8.	Karabegov quantization	16
4.9.	Karabegov's formal Berezin transform	16
5.	The disc bundle and global operators	17
5.1.	The disc bundle	17
5.2.	The generalized Hardy space	18
5.3.	The Toeplitz structure	18
5.4.	A sketch of the proof of Theorem 3.3	19
6.	Coherent States and Berezin symbols	20
6.1.	Coherent States	20
6.2.	Covariant Berezin symbols	21
6.3.	Rawnsley's ϵ function	21
6.4.	Contravariant Berezin symbols	23
6.5.	Berezin star product	24
7.	Berezin transform	25
7.1.	The definition	25
7.2.	The asymptotic expansion	26
7.3.	Norm preservation of the BT operators	26
7.4.	Bergman kernel	27
7.5.	Identification of the BT star product	28
7.6.	Pullback of the Fubini-Study form	28
	References	29

1. INTRODUCTION

For quantizable Kähler manifolds the Berezin - Toeplitz (BT) quantization scheme, both the operator quantization and the deformation quantization, supplies canonically defined quantizations. Some time ago, in joint work with Martin Bordemann and Eckhard Meinrenken, the author of this review showed that for compact Kähler manifolds it is a well-defined quantization scheme with correct semi-classical limit [21].

What makes the Berezin-Toeplitz quantization scheme so attractive is that it does not depend on further choices and that it does not only produce a formal deformation quantization, but one which is deeply related to some operator calculus.

From the point of view of classical mechanics compact Kähler manifolds appear as phase space manifolds of restricted systems or of reduced systems. A typical example of its appearance is given by the spherical pendulum which after reduction has as phase-space the complex projective space.

Very recently, inspired by fruitful applications of the basic techniques of the Berezin-Toeplitz scheme beyond the quantization of classical systems, the interest in it revived considerably.

For example these techniques show up in non-commutative geometry. More precisely, they appear in the approach to non-commutative geometry using fuzzy manifolds. The

quantum spaces of the BT quantization of level m , defined in Section 3 further down, are finite-dimensional and the quantum operator of level m constitute finite-dimensional non-commutative matrix algebras. This is the arena of non-commutative fuzzy manifolds and gauge theories over them. The classical limit, the commutative manifold, is obtained as limit $m \rightarrow \infty$. The name *fuzzy sphere* was coined by John Madore [65] for a certain quantized version of the Riemann sphere. It turned out to be a quite productive direction in the non-commutative geometry approach to quantum field theory. It is impossible to give a rather complete list of people working in this approach. The following is a rather erratic and random choice of references [12], [13], [62], [38], [11], [46], [47], [33].

Another appearance of Berezin-Toeplitz quantization techniques as basic ingredients is in the pioneering work of Jørgen Andersen on the mapping class group (MCG) of surfaces in the context of Topological Quantum Field Theory (TQFT). Beside other results, he was able to prove the asymptotic faithfulness of the mapping group action on the space of covariantly constant sections of the Verlinde bundle with respect to the Axelrod-Witten-de la Pietra and Witten connection [5, 6], see also [80]. Furthermore, he showed that the MCG does not have Kazhdan's property T . Roughly speaking, a group has *property T* says that the identity representation is isolated in the space of all unitary representations of the group [7]. In these applications the manifolds to be quantized are the moduli spaces of certain flat connections on Riemann surfaces or, equivalently, the moduli space of stable algebraic vector bundles over smooth projective curves. Here further exciting research is going on. In particular, in the realm of TQFT and the construction of modular functors [8], [9, 10].

In general quite often moduli spaces come with a Kähler structure which is quantizable. Hence, it is not surprising that the Berezin-Toeplitz quantization scheme is of importance in moduli space problems. Non-commutative deformations, and a quantization is a non-commutative deformation, yield also informations about the commutative situation. These aspects clearly need further investigations.

There are a lot of other applications on which work has already been done, recently started, or can be expected. As the Berezin - Toeplitz scheme has become a basic tool it seems the right time to collect the techniques and results in such a review. We deliberately concentrate on the case of compact Kähler manifolds. In particular, we stress the methods and results valid for all of them. Due to "space-time" limitations we will not deal with the non-compact situation. In this situation case by case studies of the examples, or class of examples are needed. See Section 3.7 for references to some of them. Also we have to skip presenting recent attempts to deal with special singular situations, like orbifolds, but see at least [63, 64], [31].

Of course, there are other reviews presenting similar quantization schemes. A very incomplete list is the following [3], [4], [2], [82], [37].

This review is self-contained in the following sense. I try to explain all notions and concepts needed to understand the results and theorems only assuming some background in modern geometry and analysis. And as such it should be accessible for a newcomer to the field (both for mathematicians as for physicists) and help him to enter these interesting research directions. It is not self-contained in the strict sense as it does supply only those proofs or sketches of proofs which are either not available elsewhere or are essential for the understanding of the

statements and concepts. The review does not require a background in quantum physics as only mathematical aspects of quantizations are touched on.

2. THE SET-UP OF GEOMETRIC QUANTIZATION

In the following I will recall the principal set-up of geometric quantization which is usually done for symplectic manifolds in the case when the manifold is a Kähler manifold.

2.1. Kähler manifolds. We will only consider phase-space manifolds which carry the structure of a Kähler manifold (M, ω) . Recall that in this case M is a complex manifold and ω , the Kähler form, is a non-degenerate closed positive $(1, 1)$ -form.

If the complex dimension of M is n then the Kähler form ω can be written with respect to local holomorphic coordinates $\{z_i\}_{i=1, \dots, n}$ as

$$\omega = i \sum_{i,j=1}^n g_{ij}(z) dz_i \wedge d\bar{z}_j, \quad (2.1)$$

with local functions $g_{ij}(z)$ such that the matrix $(g_{ij}(z))_{i,j=1, \dots, n}$ is hermitian and positive definite.

Later on we will assume that M is a compact Kähler manifold.

2.2. Poisson algebra. Denote by $C^\infty(M)$ the algebra of complex-valued (arbitrary often) differentiable functions with the point-wise multiplication as associative product. A symplectic form on a differentiable manifold is a closed non-degenerate 2-form. In particular, we can consider our Kähler form ω as a symplectic form.

For symplectic manifolds we can introduce on $C^\infty(M)$ a Lie algebra structure, the Poisson bracket *Poisson bracket* $\{.,.\}$, in the following way. First we assign to every $f \in C^\infty(M)$ its *Hamiltonian vector field* X_f , and then to every pair of functions f and g the *Poisson bracket* $\{.,.\}$ via

$$\omega(X_f, \cdot) = df(\cdot), \quad \{f, g\} := \omega(X_f, X_g). \quad (2.2)$$

One verifies that this is indeed a Lie algebra and that furthermore we have the Leibniz rule

$$\{fg, h\} = f\{g, h\} + \{f, h\}g, \quad \forall f, g, h \in C^\infty(M).$$

Such a compatible structure is called a *Poisson algebra*.

2.3. Quantum line bundles. A *quantum line bundle* for a given symplectic manifold (M, ω) is a triple (L, h, ∇) , where L is a complex line bundle, h a Hermitian metric on L , and ∇ a connection compatible with the metric h such that the (pre)quantum condition

$$\begin{aligned} \text{curv}_{L, \nabla}(X, Y) &:= \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]} = -i\omega(X, Y), \\ \text{resp. } \text{curv}_{L, \nabla} &= -i\omega \end{aligned} \quad (2.3)$$

is fulfilled. A symplectic manifold is called *quantizable* if there exists a quantum line bundle.

In the situation of Kähler manifolds we require for a quantum line bundle that it is holomorphic and that the connection is compatible both with the metric h and the complex structure of the bundle. In fact, by this requirement ∇ will be uniquely fixed. If we choose local holomorphic coordinates on the manifold and a local holomorphic frame of the bundle the

metric h will be represented by a function $\hat{h}$. In this case the curvature in the bundle can be given by $\bar{\partial}\partial \log \hat{h}$ and the quantum condition reads as

$$i \bar{\partial}\partial \log \hat{h} = \omega . \quad (2.4)$$

2.4. Example: The Riemann sphere. The Riemann sphere is the complex projective line $\mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\} \cong S^2$. With respect to the quasi-global coordinate z the form can be given as

$$\omega = \frac{i}{(1 + z\bar{z})^2} dz \wedge d\bar{z} . \quad (2.5)$$

For the Poisson bracket one obtains

$$\{f, g\} = i(1 + z\bar{z})^2 \left(\frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial g}{\partial z} - \frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} \right) . \quad (2.6)$$

Recall that the points in $\mathbb{P}^1(\mathbb{C})$ correspond to lines in $\mathbb{C}^2$ passing through the origin. If we assign to every point in $\mathbb{P}^1(\mathbb{C})$ the line it represents we obtain a holomorphic line bundle, called the tautological line bundle. The hyper plane section bundle is dual to the tautological bundle. It turns out that it is a quantum line bundle. Hence $\mathbb{P}^1(\mathbb{C})$ is quantizable.

2.5. Example: The complex projective space. Next we consider the n -dimensional complex projective space $\mathbb{P}^n(\mathbb{C})$. The example above can be extended to the projective space of any dimension. The Kähler form is given by the Fubini-Study form

$$\omega_{FS} := i \frac{(1 + |w|^2) \sum_{i=1}^n dw_i \wedge d\bar{w}_i - \sum_{i,j=1}^n \bar{w}_i w_j dw_i \wedge d\bar{w}_j}{(1 + |w|^2)^2} . \quad (2.7)$$

The coordinates w_j , $j = 1, \dots, n$ are affine coordinates $w_j = z_j/z_0$ on the affine chart $U_0 := \{(z_0 : z_1 : \dots : z_n) \mid z_0 \neq 0\}$. Again, $\mathbb{P}^n(\mathbb{C})$ is quantizable with the hyper plane section bundle as quantum line bundle.

2.6. Example: The torus. The (complex-) one-dimensional torus can be given as $M = \mathbb{C}/\Gamma_\tau$ where $\Gamma_\tau := \{n + m\tau \mid n, m \in \mathbb{Z}\}$ is a lattice with $\text{Im } \tau > 0$. As Kähler form we take

$$\omega = \frac{i\pi}{\text{Im } \tau} dz \wedge d\bar{z} , \quad (2.8)$$

with respect to the coordinate z on the covering space $\mathbb{C}$. Clearly this form is invariant under the lattice Γ_τ and hence well-defined on M . For the Poisson bracket one obtains

$$\{f, g\} = i \frac{\text{Im } \tau}{\pi} \left(\frac{\partial f}{\partial \bar{z}} \cdot \frac{\partial g}{\partial z} - \frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} \right) . \quad (2.9)$$

The corresponding quantum line bundle is the theta line bundle of degree 1, i.e. the bundle whose global sections are scalar multiples of the Riemann theta function.

2.7. Example: The unit disc and Riemann surfaces. The unit disc

$$\mathcal{D} := \{z \in \mathbb{C} \mid |z| < 1\} \quad (2.10)$$

is a non-compact Kähler manifold. The Kähler form is given by

$$\omega = \frac{2i}{(1 - z\bar{z})^2} dz \wedge d\bar{z} . \quad (2.11)$$

Every compact Riemann surface M of genus $g \geq 2$ can be given as the quotient of the unit disc under the fractional linear transformations of a Fuchsian subgroup of $SU(1,1)$. If

$R = \begin{pmatrix} a & b \\ \bar{b} & \bar{a} \end{pmatrix}$ with $|a|^2 - |b|^2 = 1$ (as an element of $SU(1,1)$) then the action is

$$z \mapsto R(z) := \frac{az + b}{\bar{b}z + \bar{a}} . \quad (2.12)$$

The Kähler form (2.11) is invariant under the fractional linear transformations. Hence it defines a Kähler form on M . The quantum bundle is the canonical bundle, i.e. the bundle whose local sections are the holomorphic differentials. Its global sections can be identified with the automorphic forms of weight 2 with respect to the Fuchsian group.

2.8. Consequences of quantizability. The above examples might create the wrong impression that every Kähler manifold is quantizable. This is not the case. For example only those higher dimensional tori complex tori are quantizable which are abelian varieties, i.e. which admit enough theta functions. It is well-known that for $n \geq 2$ a generic torus will not be an abelian variety. Why this implies that they will not be quantizable we will see in a moment.

In the language of differential geometry a line bundle is called a positive line bundle if its curvature form (up to a factor of $1/i$) is a positive form. As the Kähler form is positive the quantum condition (2.3) yields that a quantum line bundle L is a positive line bundle.

2.9. Embedding into projective space. In the following we assume that M is a quantizable compact Kähler manifold with quantum line bundle L . Kodaira's embedding theorem says that L is ample, i.e. that there exists a certain tensor power L^{m_0} of L such that the global holomorphic sections of L^{m_0} can be used to embed the phase space manifold M into the projective space of suitable dimension. The embedding is defined as follows. Let $\Gamma_{hol}(M, L^{m_0})$ be the vector space of global holomorphic sections of the bundle L^{m_0} . Fix a basis $s_0, s_1, \dots, s_N$. We choose local holomorphic coordinates z for M and a local holomorphic frame $e(z)$ for the bundle L . After these choices the basis elements can be uniquely described by local holomorphic functions $\hat{s}_0, \hat{s}_1, \dots, \hat{s}_N$ defined via $s_j(z) = \hat{s}_j(z)e(z)$. The embedding is given by the map

$$M \hookrightarrow \mathbb{P}^N(\mathbb{C}), \quad z \mapsto \phi(z) = (\hat{s}_0(z) : \hat{s}_1(z) : \dots : \hat{s}_N(z)) . \quad (2.13)$$

Note that the point $\phi(z)$ in projective space neither depends on the choice of local coordinates nor on the choice of the local frame for the bundle L . Furthermore a different choice of basis correspond to a $\mathrm{PGL}(N, \mathbb{C})$ action on the embedding space and hence the embeddings are projectively equivalent.

By this embedding quantizable compact Kähler manifolds are complex submanifolds of projective spaces. By Chow's theorem [81] they can be given as zero sets of homogenous polynomials, i.e. they are smooth projective varieties. The converse is also true. Given a

smooth subvariety M of $\mathbb{P}^n(\mathbb{C})$ it will become a Kähler manifold by restricting the Fubini-Study form. The restriction of the hyper plane section bundle will be an associated quantum line bundle.

At this place a warning is necessary. the embedding is only an embedding as complex manifolds not an isometric embedding as Kähler manifolds. This means that in general $\phi^{-1}(\omega_{FS}) \neq \omega$. See Section 7.6 for results on an “asymptotic expansion” of the pullback.

A line bundles whose global holomorphic sections will define an embedding into projective space, is called a *very ample line bundle*. In the following we will assume that L is already very ample. If L is not very ample we choose $m_0 \in \mathbb{N}$ such that the bundle L^{m_0} is very ample and take this bundle as quantum line bundle with respect to the rescaled Kähler form $m_0 \omega$ on M . The underlying complex manifold structure will not change.

3. BEREZIN-TOEPLITZ OPERATORS

In this section we will consider an operator quantization. This says that we will assign to each differentiable¹ function f on our Kähler manifold M (i.e. on our “phase space”) the Berezin-Toeplitz (BT) quantum operator T_f . More precisely, we will consider a whole family of operators $T_f^{(m)}$. These operators are defined in a canonical way. As we know from the Groenewold-van Howe theorem we cannot expect that the Poisson bracket on M can be represented by the Lie algebra of operators if we require certain desirable conditions, see [1] for further details. The best we can expect is that we obtain it at least “asymptotically”. In fact, this is true.

In our context also the operator of geometric quantization exists. At the end of this section we will discuss its relation to the BT quantum operator. It will turn out that if we take for the geometric quantization the Kähler polarization then they have the same asymptotic behaviour.

3.1. Tensor powers of the quantum line bundle. Let (M, ω) be a compact quantizable Kähler manifold and (L, h, ∇) a quantum line bundle. We assume that L is already very ample. We consider all its tensor powers

$$(L^m, h^{(m)}, \nabla^{(m)}). \quad (3.1)$$

Here $L^m := L^{\otimes m}$. If $\hat{h}$ corresponds to the metric h with respect to a local holomorphic frame e of the bundle L then $\hat{h}^m$ corresponds to the metric $h^{(m)}$ with respect to the frame $e^{\otimes m}$ for the bundle L^m . The connection $\nabla^{(m)}$ will be the induced connection.

We introduce a scalar product on the space of sections. In this review we adopt the convention that a hermitian metric (and a scalar product) is anti-linear in the first argument and linear in the second argument. First we take the Liouville form $\Omega = \frac{1}{n!} \omega^{\wedge n}$ as volume form on M and then set for the scalar product and the norm

$$\langle \varphi, \psi \rangle := \int_M h^{(m)}(\varphi, \psi) \Omega, \quad \|\varphi\| := \sqrt{\langle \varphi, \varphi \rangle}, \quad (3.2)$$

on the space $\Gamma_\infty(M, L^m)$ of global C^∞ -sections. Let $L^2(M, L^m)$ be the L^2 -completion of $\Gamma_\infty(M, L^m)$, and $\Gamma_{hol}(M, L^m)$ be its (due to the compactness of M) finite-dimensional closed

¹differentiable will always mean differentiable to any order

subspace of global holomorphic sections. Let

$$\Pi^{(m)} : L^2(M, L^m) \rightarrow \Gamma_{hol}(M, L^m) \quad (3.3)$$

be the projection.

Definition 3.1. For $f \in C^\infty(M)$ the *Toeplitz operator* $T_f^{(m)}$ (of level m) is defined by

$$T_f^{(m)} := \Pi^{(m)}(f \cdot) : \Gamma_{hol}(M, L^m) \rightarrow \Gamma_{hol}(M, L^m) . \quad (3.4)$$

In words: One takes a holomorphic section s and multiplies it with the differentiable function f . The resulting section $f \cdot s$ will only be differentiable. To obtain a holomorphic section, one has to project it back on the subspace of holomorphic sections.

The linear map

$$T^{(m)} : C^\infty(M) \rightarrow \text{End}(\Gamma_{hol}(M, L^m)), \quad f \mapsto T_f^{(m)} = \Pi^{(m)}(f \cdot) , m \in \mathbb{N}_0 . \quad (3.5)$$

is the *Toeplitz* or *Berezin-Toeplitz quantization map* (of level m). It will neither be a Lie algebra homomorphism nor an associative algebra homomorphism as in general

$$T_f^{(m)} T_g^{(m)} = \Pi^{(m)}(f \cdot) \Pi^{(m)}(g \cdot) \Pi^{(m)} \neq \Pi^{(m)}(fg \cdot) \Pi = T_{fg}^{(m)} .$$

Furthermore, on a fixed level m it is a map from the infinite-dimensional commutative algebra of functions to a noncommutative finite-dimensional (matrix) algebra. The finite-dimensionality is due to the compactness of M . A lot of classical information will get lost. To recover this information one has to consider not just a single level m but all levels together.

Definition 3.2. The Berezin-Toeplitz quantization is the map

$$C^\infty(M) \rightarrow \prod_{m \in \mathbb{N}_0} \text{End}(\Gamma_{hol}(M, L^m)), \quad f \mapsto (T_f^{(m)})_{m \in \mathbb{N}_0} . \quad (3.6)$$

We obtain a family of finite-dimensional(matrix) algebras and a family of maps. This infinite family should in some sense “approximate” the algebra $C^\infty(M)$.

3.2. Approximation results. Denote for $f \in C^\infty(M)$ by $|f|_\infty$ the sup-norm of f on M and by

$$\|T_f^{(m)}\| := \sup_{\substack{s \in \Gamma_{hol}(M, L^m) \\ s \neq 0}} \frac{\|T_f^{(m)} s\|}{\|s\|} \quad (3.7)$$

the operator norm with respect to the norm (3.2) on $\Gamma_{hol}(M, L^m)$. The following theorem was shown in 1994.

Theorem 3.3. [Bordemann, Meinrenken, Schlichenmaier] [21]

(a) For every $f \in C^\infty(M)$ there exists a $C > 0$ such that

$$|f|_\infty - \frac{C}{m} \leq \|T_f^{(m)}\| \leq |f|_\infty . \quad (3.8)$$

In particular, $\lim_{m \rightarrow \infty} \|T_f^{(m)}\| = |f|_\infty$.

(b) For every $f, g \in C^\infty(M)$

$$\|m i [T_f^{(m)}, T_g^{(m)}] - T_{\{f, g\}}^{(m)}\| = O\left(\frac{1}{m}\right) . \quad (3.9)$$

(c) For every $f, g \in C^\infty(M)$

$$\|T_f^{(m)}T_g^{(m)} - T_{f \cdot g}^{(m)}\| = O\left(\frac{1}{m}\right). \quad (3.10)$$

These results are contained in Theorem 4.1, 4.2, and in Section 5 in [21]. We will indicate the proof for (b) and (c) in Section 5. It will make reference to the symbol calculus of generalised Toeplitz operators as developed by Boutet-de-Monvel and Guillemin [24]. The original proof of (a) was quite involved and required Hermite distributions and related objects. On the basis of the asymptotic expansion of the Berezin transform [58] a more direct proof can be given. I will discuss this in Section 7.3.

Only on the basis of this theorem we are allowed to call our scheme a quantizing scheme. The properties in the theorem might be rephrased as *the BT operator quantization has the correct semiclassical limit*.

3.3. Further properties. From Theorem 3.3 (c) the

Proposition 3.4. *Let $f_1, f_2, \dots, f_r \in C^\infty(M)$ then*

$$\|T_{f_1 \dots f_r}^{(m)} - T_{f_1}^{(m)} \dots T_{f_r}^{(m)}\| = O(m^{-1}) \quad (3.11)$$

follows directly.

Proposition 3.5.

$$\lim_{m \rightarrow \infty} \| [T_f^{(m)}, T_g^{(m)}] \| = 0. \quad (3.12)$$

Proof. Using the left side of the triangle inequality, from Theorem 3.3 (b) it follows that

$$\left| m \| [T_f^{(m)}, T_g^{(m)}] \| - \| T_{\{f,g\}}^{(m)} \| \right| \leq \| m i [T_f^{(m)}, T_g^{(m)}] - T_{\{f,g\}}^{(m)} \| = O\left(\frac{1}{m}\right).$$

By part (a) of the theorem $\|T_{\{f,g\}}^{(m)}\| \rightarrow \|\{f, g\}\|_\infty$, and it stays finite. Hence $\| [T_f^{(m)}, T_g^{(m)}] \|$ has to be a zero sequence. $\square$

Proposition 3.6. *The Toeplitz map*

$$C^\infty(M) \rightarrow \text{End}(\Gamma_{\text{hol}}(M, L^{(m)})), \quad f \mapsto T_f^{(m)},$$

is surjective.

For a proof see [21, Prop. 4.2].

This proposition says that for a fixed m every operator $A \in \text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ is the Toeplitz operator of a function f_m . In the language of Berezin's co- and contravariant symbols f_m will be the contravariant symbol of A . We will discuss this in Section 6.2.

Proposition 3.7. *For all $f \in C^\infty(M)$*

$$T_f^{(m)*} = T_{\bar{f}}^{(m)}.$$

In particular, for real valued functions f the associated Toeplitz operator is selfadjoint.

Proof. Take $s, t \in \Gamma_{\text{hol}}(M, L^m)$ then

$$\langle s, T_f^{(m)} t \rangle = \langle s, \Pi^{(m)}(f \cdot t) \rangle = \langle s, f \cdot t \rangle = \langle \bar{f} \cdot s, t \rangle = \langle T_{\bar{f}}^{(m)} s, t \rangle.$$

$\square$

The opposite of the last statement of the above proposition is also true in the following sense.

Proposition 3.8. *Let $A \in \text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ be a selfadjoint operator then there exists a real valued function f , such that $A = T_f^{(m)}$.*

Proof. By the surjectivity of the Toeplitz map $A = T_f^{(m)}$ with a complex valued function $f = f_0 + i f_1$ with real functions f_0 and f_1 . As $T_f^{(m)} = A = A^* = T_{\bar{f}}^{(m)}$ it follows $T_{f-\bar{f}} = 0$ and hence $T_{f_1}^{(m)} = 0$. From this we conclude $A = T_f^{(m)} = T_{f_0}^{(m)}$. $\square$

We like to stress the fact that the Toeplitz map is never injective on a fixed level m . Only if $\|T_{f-g}^{(m)}\| \rightarrow 0$ for $m \rightarrow 0$ we can conclude that $f = g$.

Proposition 3.9. *Let $f \in C^\infty(M)$ and $n = \dim_{\mathbb{C}} M$. Denote the trace on $\text{End}(\Gamma_{\text{hol}}(M, L^m))$ by $\text{Tr}^{(m)}$ then*

$$\text{Tr}^{(m)}(T_f^{(m)}) = m^n \left(\frac{1}{\text{vol}(\mathbb{P}^n(\mathbb{C}))} \int_M f \Omega + O(m^{-1}) \right). \quad (3.13)$$

See [21], resp. [78] for a detailed proof.

3.4. Strict quantization. The asymptotic results of Theorem 3.3 says that the BT operator quantization is a strict quantization in the sense of Rieffel [73] as formulated in the book by Landsman [61]. We take as base space $X = \{0\} \cup \{1/m \mid m \in \mathbb{N}\}$, with its induced topology coming from $\mathbb{R}$. Note that $\{0\}$ is an accumulation point of the set $\{1/m \mid m \in \mathbb{N}\}$. As C^* algebras above the points $\{1/m\}$ we take the algebras $\text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ and above $\{0\}$ the algebra $C^\infty(M)$. For $f \in C^\infty(M)$ we assign $0 \mapsto f$, and $1/m \mapsto T_f^{(m)}$. Now the property (a) in Theorem 3.3 is called in [61] Rieffel's condition, (b) Dirac's condition, and (c) von Neumann's condition. Completeness is true by Propositions 3.6 and 3.8.

This definition is closely related to the notion of continuous fields of C^* -algebras, see [61].

3.5. Relation to geometric quantization. There exists another quantum operator in the geometric setting, the operator of geometric quantization introduced by Kostant and Souriau. In a first step the prequantum operator associated to the bundle L^m for the function $f \in C^\infty(M)$ is defined as

$$P_f^{(m)} := \nabla_{X_f^{(m)}}^{(m)} + i f \cdot \text{id}. \quad (3.14)$$

Here $\nabla^{(m)}$ is the connection in L^m , and $X_f^{(m)}$ the Hamiltonian vector field of f with respect to the Kähler form $\omega^{(m)} = m \cdot \omega$, i.e. $m\omega(X_f^{(m)}, \cdot) = df(\cdot)$. This operator $P_f^{(m)}$ acts on the space of differentiable global sections of the line bundle L^m . The sections depend at every point on $2n$ local coordinates and one has to restrict the space to sections covariantly constant along the excessive dimensions. In technical terms, one chooses a *polarization*. In general such a polarization is not unique. But in our complex situation there is canonical one by only taking the holomorphic sections. This polarization is called *Kähler polarization*. The operator of geometric quantization is then defined by

$$Q_f^{(m)} := \Pi^{(m)} P_f^{(m)}. \quad (3.15)$$

The Toeplitz operator and the operator of geometric quantization (with respect to the Kähler polarization) are related by

Proposition 3.10. (*Tuynman Lemma*) *Let M be a compact quantizable Kähler manifold then*

$$Q_f^{(m)} = i \cdot T_{f - \frac{1}{2m}\Delta f}^{(m)}, \quad (3.16)$$

where Δ is the Laplacian with respect to the Kähler metric given by ω .

For the proof see [84], and [20] for a coordinate independent proof.

In particular the $Q_f^{(m)}$ and the $T_f^{(m)}$ have the same asymptotic behaviour. We obtain for $Q_f^{(m)}$ similar results as in Theorem 3.3. For details see [74]. It should be noted that for (3.16) the compactness of M is essential.

3.6. L_α approximation. In [20] the notion of L_α , resp. $gl(N)$, resp. $su(N)$ quasi-limit were introduced. It was conjectured in [20] that for every compact quantizable Kähler manifold the Poisson algebra of functions is a $gl(N)$ quasi-limit. In fact, the conjecture follows from the Theorem 3.3, see [21] and [74] for details.

3.7. The noncompact situation. Berezin-Toeplitz operators can be introduced for non-compact Kähler manifolds. In this case the L^2 spaces are the space of bounded sections and for the subspaces of holomorphic sections one can only consider the bounded holomorphic sections. Unfortunately, in this context the proofs of Theorem 3.3 do not work. One has to study examples or classes of examples case by case whether the corresponding properties are correct.

In the following we give a very incomplete list of references. Berezin himself studied bounded complex-symmetric domains [17]. In this case the manifold is an open domain in $\mathbb{C}^n$. Instead of sections one studies functions which are integrable with respect to a suitable measure depending on $\hbar$. Then $1/\hbar$ corresponds to the tensor power of our bundle. Such Toeplitz operators were studied extensively by Upmeyer in a series of works [87, 86, 88, 89]. See also the book of Upmeyer [90]. For $\mathbb{C}^n$ see Berger and Coburn [18], [34]. Klimek and Lesniewski [59] studied the Berezin-Toeplitz quantization on the unit disc. Using automorphic forms and the universal covering they obtain results for Riemann surfaces of genus $g \geq 2$. The names of Borthwick, Klimek, Lesniewski, Rinaldi, and Upmeyer should be mentioned in the context of BT quantization for Cartan domains and super Hermitian spaces.

A quite different approach to Berezin-Toeplitz quantization is based on the asymptotic expansion of the Bergman kernel outside the diagonal. This was also used by the author together with Karabegov [58] for the compact Kähler case. See Section 7 for some details. Engliš [42] showed similar results for bounded pseudo-convex domains in $\mathbb{C}^N$. Ma and Marinescu [63, 64] developed a theory of Bergman kernels for the symplectic case, which yields also results on the Berezin-Toeplitz operators for certain non-compact Kähler manifolds and even orbifolds.

4. BEREZIN-TOEPLITZ DEFORMATION QUANTIZATION

There is another approach to quantization. Instead of assigning noncommutative operators to commuting functions one might think about “deforming” the pointwise commutative multiplication of functions into a non-commutative product. It is required to remain associative,

the commutator of two elements should relate to the Poisson bracket of the elements, and it should reduce in the “classical limit” to the commutative situation.

It turns out that such a deformation which is valid for all differentiable functions cannot exist. A way out is to enlarge the algebra of functions by considering formal power series over them and to deform the product inside this bigger algebra. A first systematic treatment and applications in physics of this idea were given 1978 by Bayen, Flato, Fronsdal, Lichnerowicz, and Sternheimer [14]. There the notion of *deformation quantization* and *star products* were introduced. Earlier versions of these concepts were around due to Berezin [16], Moyal [68], and Weyl [91]. For a presentation of the history see [82].

We will show that for compact Kähler manifolds M , there is a natural star product.

4.1. Definition of star products. We start with a Poisson manifold $(M, \{.,.\})$, i.e. a differentiable manifold with a Poisson bracket for the function such that $(C^\infty(M), \cdot, \{.,.\})$ is a Poisson algebra. Let $\mathcal{A} = C^\infty(M)[[\nu]]$ be the algebra of formal power series in the variable ν over the algebra $C^\infty(M)$.

Definition 4.1. A product $\star$ on $\mathcal{A}$ is called a (formal) star product for M (or for $C^\infty(M)$) if it is an associative $\mathbb{C}[[\nu]]$ -linear product which is ν -adically continuous such that

- (1) $\mathcal{A}/\nu\mathcal{A} \cong C^\infty(M)$, i.e. $f \star g \bmod \nu = f \cdot g$,
- (2) $\frac{1}{\nu}(f \star g - g \star f) \bmod \nu = -i\{f, g\}$,

where $f, g \in C^\infty(M)$.

Alternatively we can write

$$f \star g = \sum_{j=0}^{\infty} C_j(f, g) \nu^j, \quad (4.1)$$

with $C_j(f, g) \in C^\infty(M)$ such that the C_j are bilinear in the entries f and g . The conditions (1) and (2) can be reformulated as

$$C_0(f, g) = f \cdot g, \quad \text{and} \quad C_1(f, g) - C_1(g, f) = -i\{f, g\}. \quad (4.2)$$

By the ν -adic continuity (4.1) fixes $\star$ on $\mathcal{A}$. A (formal) deformation quantization is given by a (formal) star product. I will use both terms interchangeable.

There are certain additional conditions for a star product which are sometimes useful.

- (1) We call it “null on constants”, if $1 \star f = f \star 1 = f$, which is equivalent to the fact that the constant function 1 will remain the unit in $\mathcal{A}$. In terms of the coefficients it can be formulated as $C_k(f, 1) = C_k(1, f) = 0$ for $k \geq 1$. In this review we always assume this to be the case for star products.
- (2) We call it selfadjoint if $\overline{f \star g} = \overline{g} \star \overline{f}$, where we assume $\bar{\nu} = \nu$.
- (3) We call it local if

$$\text{supp } C_j(f, g) \subseteq \text{supp } f \cap \text{supp } g, \quad \forall f, g \in C^\infty(M).$$

From the locality property it follows that the C_j are bidifferential operators and that the global star product defines for every open subset U of M a star product for the Poisson algebra $C^\infty(U)$. Such local star products are also called *differential star products*.

4.2. Existence of star products. In the usual setting of deformation theory there always exists a trivial deformation. This is not the case here, as the trivial deformation of $C^\infty(M)$ to $\mathcal{A}$, which is nothing else as extending the point-wise product to the power series, is not allowed as it does not fulfil Condition (2) in Definition 4.1 (at least not if the Poisson bracket is non-trivial). In fact the existence problem is highly non-trivial. In the symplectic case different existence proofs, from different perspectives, were given by DeWilde-Lecomte [35], Omori-Maeda-Yoshioka [70], and Fedosov [45]. The general Poisson case was settled by Kontsevich [60].

4.3. Equivalence and classification of star products.

Definition 4.2. Given a Poisson manifold $(M, \{.,.\})$. Two star products $\star$ and $\star'$ associated to the Poisson structure $\{.,.\}$ are called equivalent if and only if there exists a formal series of linear operators

$$B = \sum_{i=0}^{\infty} B_i \nu^i, \quad B_i : C^\infty(M) \rightarrow C^\infty(M), \quad (4.3)$$

with $B_0 = id$ such that

$$B(f) \star' B(g) = B(f \star g). \quad (4.4)$$

For local star products in the general Poisson setting there are complete classification results. Here I will only consider the symplectic case.

To each local star product $\star$ its *Fedosov-Deligne class*

$$cl(\star) \in \frac{1}{i\nu}[\omega] + H_{dR}^2(M)[[\nu]] \quad (4.5)$$

can be assigned. Here $H_{dR}^2(M)$ denotes the 2nd deRham cohomology class of closed 2-forms modulo exact forms and $H_{dR}^2(M)[[\nu]]$ the formal power series with such classes as coefficients. Such formal power series are called *formal deRham classes*. In general we will use $[\alpha]$ for the cohomology class of a form α .

This assignment gives a 1:1 correspondence between the formal deRham classes and the equivalence classes of star products.

For contractible manifolds we have $H_{dR}^2(M) = 0$ and hence there is up to equivalence exactly one local star product. This yields that locally all local star products of a manifold are equivalent to a certain fixed one, which is called the Moyal product. For these and related classification results see [36], [50], [19], [69].

4.4. Star products with separation of variables. For our compact Kähler manifolds we will have many different and even non-equivalent star products. The question is: is there a star product which is given in a natural way? The answer will be yes: the Berezin-Toeplitz star product to be introduced below. First we consider star products respecting the complex structure in a certain sense.

Definition 4.3. (Karabegov [53]) A star product is called *star product with separation of variables* if and only if

$$f \star h = f \cdot h, \quad \text{and} \quad h \star g = h \cdot g, \quad (4.6)$$

for every locally defined holomorphic function g , antiholomorphic function f , and arbitrary function h .

Recall that a local star product $\star$ for M defines a star product for every open subset U of M . We have just to take the bidifferential operators defining $\star$. Hence it makes sense to talk about $\star$ -multiplying with local functions.

Proposition 4.4. *A local $\star$ product has the separation of variables property if and only if in the bidifferential operators $C_k(.,.)$ for $k \geq 1$ in the first argument only derivatives in holomorphic and in the second argument only derivatives in antiholomorphic directions appear.*

In Karabegov's original notation the rôles of the holomorphic and antiholomorphic functions is switched. Bordemann and Waldmann [22] called such star products *star products of Wick type*. Both Karabegov and Bordemann-Waldmann proved that there exists for every Kähler manifold star products of separation of variables type. In Section 4.8 we will give more details on Karabegov's construction. Bordemann and Waldmann modified Fedosov's method [45] to obtain such a star product. See also Reshetikhin and Takhtajan [72] for yet another construction. But I like to point out that in all these constructions the result is only a formal star product without any relation to an operator calculus, which will be given by the Berezin-Toeplitz star product introduced in the next section.

Another warning is in order. The property of being a star product of separation of variables type will not be kept by equivalence transformations.

4.5. Berezin-Toeplitz star product.

Theorem 4.5. *There exists a unique (formal) star product $\star_{BT}$ for M*

$$f \star_{BT} g := \sum_{j=0}^{\infty} \nu^j C_j(f, g), \quad C_j(f, g) \in C^\infty(M), \quad (4.7)$$

in such a way that for $f, g \in C^\infty(M)$ and for every $N \in \mathbb{N}$ we have with suitable constants $K_N(f, g)$ for all m

$$\|T_f^{(m)} T_g^{(m)} - \sum_{0 \leq j < N} \left(\frac{1}{m}\right)^j T_{C_j(f, g)}^{(m)}\| \leq K_N(f, g) \left(\frac{1}{m}\right)^N. \quad (4.8)$$

The star product is null on constants and selfadjoint.

This theorem has been proven immediately after [21] was finished. It has been announced in [75], [76] and the proof was written up in German in [74]. A complete proof published in English can be found in [78].

For simplicity we might write

$$T_f^{(m)} \cdot T_g^{(m)} \sim \sum_{j=0}^{\infty} \left(\frac{1}{m}\right)^j T_{C_j(f, g)}^{(m)} \quad (m \rightarrow \infty), \quad (4.9)$$

but we will always assume the strong and precise statement of (4.8). The same is assumed for other asymptotic formulas appearing further down in this review.

Next we want to identify this star product. Let K_M be the canonical line bundle of M , i.e. the n^{th} exterior power of the holomorphic 1-differentials. The canonical class δ is the first Chern class of this line bundle, i.e. $\delta := c_1(K_M)$. If we take in K_M the fibre metric coming from the Liouville form Ω then this defines a unique connection and further a unique curvature $(1, 1)$ -form ω_{can} . In our sign conventions we have $\delta = [\omega_{can}]$.

Together with Karabegov the author showed

Theorem 4.6. [58] (a) *The Berezin-Toeplitz star product is a local star product which is of separation of variable type.*
 (b) *Its classifying Deligne-Fedosov class is*

$$cl(\star_{BT}) = \frac{1}{i} \left(\frac{1}{\nu} [\omega] - \frac{\delta}{2} \right) \quad (4.10)$$

for the characteristic class of the star product $\star_{BT}$.

(c) *The classifying Karabegov form associated to the Berezin-Toeplitz star product is*

$$-\frac{1}{\nu} \omega + \omega_{can}. \quad (4.11)$$

The Karabegov form has not yet defined here. We will introduce it below in Section 4.8. Using K -theoretic methods the formula for $cl(\star_{BT})$ was also given by Hawkins [51].

4.6. Star product of geometric quantization. Tuynman's result (3.16) relates the operators of geometric quantization with Kähler polarization and the BT operators. As the latter define a star product it can be used to give also a star product $\star_{GQ}$ associated to geometric quantization. Details can be found in [78]. This star product will be equivalent to the BT star product, but it is not of separation of variables type. The equivalence is given by the $\mathbb{C}[[\nu]]$ -linear map induced by

$$B(f) := f - \nu \frac{\Delta}{2} f = (id - \nu \frac{\Delta}{2}) f. \quad (4.12)$$

We obtain $B(f) \star_{BT} B(g) = B(f \star_{GQ} g)$.

4.7. Trace for the BT star product. From (3.13) the following complete asymptotic expansion for $m \rightarrow \infty$ can be deduced [78], [23]):

$$\mathrm{Tr}^{(m)}(T_f^{(m)}) \sim m^n \left(\sum_{j=0}^{\infty} \left(\frac{1}{m} \right)^j \tau_j(f) \right), \quad \text{with } \tau_j(f) \in \mathbb{C}. \quad (4.13)$$

We define the $\mathbb{C}[[\nu]]$ -linear map

$$\mathrm{Tr} : C^\infty(M)[[\nu]] \rightarrow \nu^{-n} \mathbb{C}[[\nu]], \quad \mathrm{Tr} f := \nu^{-n} \sum_{j=0}^{\infty} \nu^j \tau_j(f), \quad (4.14)$$

where the $\tau_j(f)$ are given by the asymptotic expansion (4.13) for $f \in C^\infty(M)$ and for arbitrary elements by $\mathbb{C}[[\nu]]$ -linear extension.

Proposition 4.7. [78] *The map Tr is a trace, i.e., we have*

$$\mathrm{Tr}(f \star g) = \mathrm{Tr}(g \star f). \quad (4.15)$$

4.8. Karabegov quantization. In [53, 54] Karabegov not only gave the notion of *separation of variables type*, but also a proof of existence of such formal star products for any Kähler manifold, whether compact, non-compact, quantizable, or non-quantizable. Moreover, he classified them completely as individual star product not only up to equivalence.

He starts with (M, ω_{-1}) a pseudo-Kähler manifold, i.e. a complex manifold with a non-degenerate closed $(1, 1)$ -form not necessarily positive.

A formal form $\widehat{\omega} = (1/\nu)\omega_{-1} + \omega_0 + \nu\omega_1 + \dots$ is called a formal deformation of the form $(1/\nu)\omega_{-1}$ if the forms ω_r , $r \geq 0$, are closed but not necessarily nondegenerate $(1, 1)$ -forms on M . It was shown in [53] that all deformation quantizations with separation of variables on the pseudo-Kähler manifold (M, ω_{-1}) are bijectively parametrized by the formal deformations of the form $(1/\nu)\omega_{-1}$.

Assume that we have such a star product $(\mathcal{A} := C^\infty(M)[[\nu]], \star)$. Then for $f, g \in \mathcal{A}$ the operators of left and right multiplication L_f, R_g are given by $L_f g = f \star g = R_g f$. The associativity of the star-product $\star$ is equivalent to the fact that L_f commutes with R_g for all $f, g \in \mathcal{A}$. If a star product is differential then L_f, R_g are formal differential operators.

Karabegov constructs his star product associated to the deformation $\widehat{\omega}$ in the following way. First he chooses on every contractible coordinate chart $U \subset M$ (with holomorphic coordinates $\{z_k\}$) its formal potential

$$\widehat{\Phi} = (1/\nu)\Phi_{-1} + \Phi_0 + \nu\Phi_1 + \dots, \quad \widehat{\omega} = i\partial\bar{\partial}\widehat{\Phi}. \quad (4.16)$$

Then construction is done in such a way that we have for the left (right) multiplication operators on U

$$L_{\partial\Phi/\partial z_k} = \partial\Phi/\partial z_k + \partial/\partial z_k, \quad \text{and} \quad R_{\partial\Phi/\partial \bar{z}_l} = \partial\Phi/\partial \bar{z}_l + \partial/\partial \bar{z}_l. \quad (4.17)$$

The set $\mathcal{L}(U)$ of all left multiplication operators on U is completely described as the set of all formal differential operators commuting with the point-wise multiplication operators by antiholomorphic coordinates $R_{\bar{z}_l} = \bar{z}_l$ and the operators $R_{\partial\Phi/\partial \bar{z}_l}$. From the knowledge of $\mathcal{L}(U)$ the star product on U can be reconstructed. The local star-products agree on the intersections of the charts and define the global star-product $\star$ on M .

We have to mention that this original construction of Karabegov will yield a star product of separation of variable type but with the role of holomorphic and antiholomorphic variables switched. This says for any open subset $U \subset M$ and any holomorphic function a and antiholomorphic function b on U the operators L_a and R_b are the operators of point-wise multiplication by a and b respectively, i.e., $L_a = a$ and $R_b = b$.

4.9. Karabegov's formal Berezin transform. Given such a star products $\star$, Karabegov introduced the formal *Berezin transform* I as the unique formal differential operator on M such that for any open subset $U \subset M$, holomorphic functions a and antiholomorphic functions b on U the relation $I(a \cdot b) = b \star a$ holds (see [55]). He shows that $I = 1 + \nu\Delta + \dots$, where Δ is the Laplace operator corresponding to the pseudo-Kähler metric on M .

Karabegov considered the following associated star products First the *dual* star-product $\tilde{\star}$ on M is defined for $f, g \in \mathcal{A}$ by the formula

$$f \tilde{\star} g = I^{-1}(Ig \star If). \quad (4.18)$$

It is a star-product with separation of variables on the pseudo-Kähler manifold $(M, -\omega_{-1})$. Its formal Berezin transform equals I^{-1} , and thus the dual to $\tilde{\star}$ is $\star$. Note that it is not a

star product of the same pseudo-Kähler manifold. Denote by $\tilde{\omega} = -(1/\nu)\omega_{-1} + \tilde{\omega}_0 + \nu\tilde{\omega}_1 + \dots$ the formal form parametrizing the star-product $\tilde{\star}$.

Next, the opposite of the dual star-product, $\star' = \tilde{\star}^{op}$, is given by the formula

$$f \star' g = I^{-1}(If \star Ig). \quad (4.19)$$

It defines a deformation quantization with separation of variables on M , but with the roles of holomorphic and antiholomorphic variables swapped - with respect to $\star$. It could be described also as a deformation quantization with separation of variables on the pseudo-Kähler manifold $(\overline{M}, \omega_{-1})$ where $\overline{M}$ is the manifold M with the opposite complex structure. But now the pseudo-Kähler form will be the same. Indeed the formal Berezin transform I establishes an equivalence of deformation quantizations $(\mathcal{A}, \star)$ and $(\mathcal{A}, \star')$.

How is the relation to the Berezin-Toeplitz star product $\star_{BT}$ of Theorem 4.5? There exists a certain formal deformation $\hat{\omega}$ of the form $(1/\nu)\omega$ which yields a star product $\star$ in the Karabegov sense. The opposite of its dual will be equal to the Berezin-Toeplitz star product, i.e.

$$\star_{BT} = \tilde{\star}^{op} = \star'. \quad (4.20)$$

The classifying Karabegov form $\tilde{\omega}$ of $\tilde{\star}$ will be the form (4.11). Note as $\star$ and $\star_{BT}$ are equivalent via I , we have $cl(\star) = cl(\star_{BT})$, see the formula (4.10). We will identify $\hat{\omega}$ in Section 7.5.

5. THE DISC BUNDLE AND GLOBAL OPERATORS

In this section we identify the bundles L^m over the Kähler manifold M as associated line bundles of one unique S^1 -bundle over M . The Toeplitz operator will appear as “modes” of a global Toeplitz operator. A detailed analysis of this global operator will yield a proof of Theorem 3.3 part (b) and part (c).

Moreover, we will need this set-up to discuss coherent states, Berezin symbols, and the Berezin transform in the next sections. For a more detailed presentation see [74].

5.1. The disc bundle. We will assume that the quantum line bundle L is already very ample, i.e. it has enough global holomorphic sections to embed M into projective space. From the bundle² (L, h) we pass to its dual $(U, k) := (L^*, h^{-1})$ with dual metric k . Inside of the total space U we consider the circle bundle

$$Q := \{\lambda \in U \mid k(\lambda, \lambda) = 1\}, \quad (5.1)$$

the (open) disc bundle and (closed) disc bundle respectively

$$D := \{\lambda \in U \mid k(\lambda, \lambda) < 1\}, \quad \overline{D} := \{\lambda \in U \mid k(\lambda, \lambda) \leq 1\}. \quad (5.2)$$

Let $\tau : U \rightarrow M$ the projection (maybe restricted to the subbundles).

For the projective space $\mathbb{P}^N(\mathbb{C})$ with the hyperplane section bundle H as quantum line bundle the bundle U is just the tautological bundle. Its fibre over the point $z \in \mathbb{P}^N(\mathbb{C})$ consists of the line in $\mathbb{C}^{N+1}$ which is represented by z . In particular, for the projective space the total space of U with the zero section removed can be identified with $\mathbb{C}^{N+1} \setminus \{0\}$. The same picture remains true for the via the very ample quantum line bundle in projective space

²As the connection ∇ will not be needed anymore, I will drop it in the notation.

embedded manifold M . The quantum line bundle will be the pull-back of H (i.e. its restriction to the embedded manifold) and its dual is the pull-back of the tautological bundle.

In the following we use $E \setminus 0$ to denote the total space of a vector bundle E with the image of the zero section removed. Starting from the real valued function $\hat{k}(\lambda) := k(\lambda, \lambda)$ on U we define $\tilde{a} := \frac{1}{2i}(\partial - \bar{\partial}) \log \hat{k}$ on $U \setminus 0$ (the derivation are taken with respect to the complex structure on U) and denote by α its restriction to Q . With the help of the quantization condition (2.3) we obtain $d\alpha = \tau^*\omega$ (with the deRham differential $d = d_Q$) and that in fact $\mu = \frac{1}{2\pi}\tau^*\Omega \wedge \alpha$ is a volume form on Q . Indeed α is a contact form for the contact manifold Q . As far as the integration is concern we get

$$\int_Q (\tau^*f)\mu = \int_M f\Omega, \quad \forall f \in C^\infty(M). \quad (5.3)$$

Recall that Ω is the Liouville volume form on M .

5.2. The generalized Hardy space. With respect to μ we take the L^2 -completion $L^2(Q, \mu)$ of the space of functions on Q . The generalized *Hardy space* $\mathcal{H}$ is the closure of the space of those functions in $L^2(Q, \mu)$ which can be extended to holomorphic functions on the whole disc bundle $\bar{D}$. The generalized *Szegő projector* is the projection

$$\Pi : L^2(Q, \mu) \rightarrow \mathcal{H}. \quad (5.4)$$

By the natural circle action the bundle Q is an S^1 -bundle and the tensor powers of U can be viewed as associated line bundles. The space $\mathcal{H}$ is preserved by the S^1 -action. It can be decomposed into eigenspaces $\mathcal{H} = \prod_{m=0}^\infty \mathcal{H}^{(m)}$ where $c \in S^1$ acts on $\mathcal{H}^{(m)}$ as multiplication by c^m . The Szegő projector is S^1 invariant and can be decomposed into its components, the Bergman projectors

$$\hat{\Pi}^{(m)} : L^2(Q, \mu) \rightarrow \mathcal{H}^{(m)}. \quad (5.5)$$

Sections of $L^m = U^{-m}$ can be identified with functions ψ on Q which satisfy the equivariance condition $\psi(c\lambda) = c^m\psi(\lambda)$, i.e. which are homogeneous of degree m . This identification is given via the map

$$\gamma_m : L^2(M, L^m) \rightarrow L^2(Q, \mu), \quad s \mapsto \psi_s \quad \text{where} \quad \psi_s(\alpha) = \alpha^{\otimes m}(s(\tau(\alpha))), \quad (5.6)$$

which turns out to be an isometry onto its image. On $L^2(M, L^m)$ we have the scalar product (3.2). Restricted to the holomorphic sections we obtain the isometry

$$\gamma_m : \Gamma_{hol}(M, L^m) \cong \mathcal{H}^{(m)}. \quad (5.7)$$

In the case of $\mathbb{P}^N(\mathbb{C})$ this correspondence is nothing else as the identification of the global sections of the m^{th} tensor powers of the hyper plane section bundle with the homogenous polynomial functions of degree m on $\mathbb{C}^{N+1}$.

5.3. The Toeplitz structure. There is the notion of Toeplitz structure (Π, Σ) as developed by Boutet de Monvel and Guillemin in [24, 48]. I do not want to present the general theory only the specialization to our situation. Here Π is the Szegő projector (5.4) and Σ is the submanifold

$$\Sigma := \{ t\alpha(\lambda) \mid \lambda \in Q, t > 0 \} \subset T^*Q \setminus 0 \quad (5.8)$$

of the tangent bundle of Q defined with the help of the 1-form α . It turns out that Σ is a symplectic submanifold, called a symplectic cone.

A (generalized) *Toeplitz operator* of order k is an operator $A : \mathcal{H} \rightarrow \mathcal{H}$ of the form $A = \Pi \cdot R \cdot \Pi$ where R is a pseudodifferential operator (Ψ DO) of order k on Q . The Toeplitz operators constitute a ring. The symbol of A is the restriction of the principal symbol of R (which lives on T^*Q) to Σ . Note that R is not fixed by A , but Guillemin and Boutet de Monvel showed that the symbols are well-defined and that they obey the same rules as the symbols of Ψ DOs. In particular, the following relations are valid:

$$\sigma(A_1 A_2) = \sigma(A_1) \sigma(A_2), \quad \sigma([A_1, A_2]) = i \{ \sigma(A_1), \sigma(A_2) \}_\Sigma. \quad (5.9)$$

Here $\{.,.\}_\Sigma$ is the restriction of the canonical Poisson structure of T^*Q to Σ coming from the canonical symplectic form on T^*Q .

5.4. A sketch of the proof of Theorem 3.3. For this we need only to consider the following two generalized Toeplitz operators:

- (1) The generator of the circle action gives the operator $D_\varphi = \frac{1}{i} \frac{\partial}{\partial \varphi}$, where φ is the angular variable. It is an operator of order 1 with symbol t . It operates on $\mathcal{H}^{(m)}$ as multiplication by m .
- (2) For $f \in C^\infty(M)$ let M_f be the operator on $L^2(Q, \mu)$ corresponding to multiplication with $\tau^* f$. We set

$$T_f = \Pi \cdot M_f \cdot \Pi : \mathcal{H} \rightarrow \mathcal{H}. \quad (5.10)$$

As M_f is constant along the fibres of τ , the operator T_f commutes with the circle action. Hence we can decompose

$$T_f = \prod_{m=0}^{\infty} T_f^{(m)}, \quad (5.11)$$

where $T_f^{(m)}$ denotes the restriction of T_f to $\mathcal{H}^{(m)}$. After the identification of $\mathcal{H}^{(m)}$ with $\Gamma_{hol}(M, L^m)$ we see that these $T_f^{(m)}$ are exactly the Toeplitz operators $T_f^{(m)}$ introduced in Section 3. We call T_f the global Toeplitz operator and the $T_f^{(m)}$ the local Toeplitz operators. The operator T_f is of order 0. Let us denote by $\tau_\Sigma : \Sigma \subseteq T^*Q \rightarrow Q \rightarrow M$ the composition then we obtain for its symbol $\sigma(T_f) = \tau_\Sigma^*(f)$.

Now we are able to proof (3.9). First we introduce for a fixed $t > 0$

$$\Sigma_t := \{t \cdot \alpha(\lambda) \mid \lambda \in Q\} \subseteq \Sigma. \quad (5.12)$$

It turns out that $\omega_{\Sigma|\Sigma_t} = -t\tau_\Sigma^*\omega$. The commutator $[T_f, T_g]$ is a Toeplitz operator of order -1 . From the above we obtain with (5.9) that the symbol of the commutator equals

$$\sigma([T_f, T_g])(t\alpha(\lambda)) = i \{ \tau_\Sigma^* f, \tau_\Sigma^* g \}_\Sigma(t\alpha(\lambda)) = -i t^{-1} \{ f, g \}_M(\tau(\lambda)). \quad (5.13)$$

We consider the Toeplitz operator

$$A := D_\varphi^2 [T_f, T_g] + i D_\varphi T_{\{f, g\}}. \quad (5.14)$$

Formally this is an operator of order 1. Using $\sigma(T_{\{f, g\}}) = \tau_\Sigma^* \{f, g\}$ and $\sigma(D_\varphi) = t$ we see that its principal symbol vanishes. Hence it is an operator of order 0. Now M and hence also Q are compact manifolds. This implies that A is a bounded operator (Ψ DOs of order 0 on compact

manifolds are bounded). It is obviously S^1 -invariant and we can write $A = \prod_{m=0}^{\infty} A^{(m)}$ where $A^{(m)}$ is the restriction of A on the space $\mathcal{H}^{(m)}$. For the norms we get $\|A^{(m)}\| \leq \|A\|$. But

$$A^{(m)} = A|_{\mathcal{H}^{(m)}} = m^2[T_f^{(m)}, T_g^{(m)}] + i m T_{\{f,g\}}^{(m)}. \quad (5.15)$$

Taking the norm bound and dividing it by m we get part (b) of Theorem 3.3. Using (5.7) the norms involved indeed coincide.

Quite similar one can prove part (c) of Theorem 3.3 and more general the existence of the coefficients $C_j(f, g)$ for the Berezin-Toeplitz star product of Theorem 4.5. See [78] and [74] for the details.

6. COHERENT STATES AND BEREZIN SYMBOLS

6.1. Coherent States. Let the situation be as in the previous section. In particular L is assumed to be already very ample, $U = L^*$ is the dual of the quantum line bundle, $Q \subset U$ the unit circle bundle, and $\tau : Q \rightarrow M$ the projection. In particular, recall the correspondence (5.6) $\psi_s(\alpha) = \alpha^{\otimes m}(s\tau(\alpha))$ of m -homogeneous functions ψ_s on U with sections of L^m . To obtain this correspondence we fixed the section s and varied α .

Now we do the opposite. We fix $\alpha \in U \setminus 0$ and vary the sections s . Obviously this yields a linear form on $\Gamma_{hol}(M, L^m)$ and hence with the help of the scalar product (3.2) we make the following

Definition 6.1. (a) The *coherent vector (of level m)* associated to the point $\alpha \in U \setminus 0$ is the unique element $e_\alpha^{(m)}$ of $\Gamma_{hol}(M, L^m)$ such that

$$\langle e_\alpha^{(m)}, s \rangle = \psi_s(\alpha) = \alpha^{\otimes m}(s(\tau(\alpha))) \quad (6.1)$$

for all $s \in \Gamma_{hol}(M, L^m)$.

(b) The *coherent state (of level m)* associated to $x \in M$ is the projective class

$$e_x^{(m)} := [e_\alpha^{(m)}] \in \mathbb{P}(\Gamma_{hol}(M, L^m)), \quad \alpha \in \tau^{-1}(x), \alpha \neq 0. \quad (6.2)$$

Of course, we have to show that the object in (b) is well-defined. Recall that $\langle \cdot, \cdot \rangle$ denotes the scalar product on the space of global sections $\Gamma_\infty(M, L^m)$. In the convention of this review it will be anti-linear in the first argument and linear in the second argument. The coherent vectors are antiholomorphic in α and fulfil

$$e_{c\alpha}^{(m)} = \bar{c}^m \cdot e_\alpha^{(m)}, \quad c \in \mathbb{C}^* := \mathbb{C} \setminus \{0\}. \quad (6.3)$$

Note that $e_\alpha^{(m)} \equiv 0$ would imply that all sections will vanish at the point $x = \tau(\alpha)$. Hence, the sections of L cannot be used to embed M into projective space, which is a contradiction to the very-ampleness of L . Hence, $e_\alpha^{(m)} \neq 0$ and due to (6.3) the class

$$[e_\alpha^{(m)}] := \{s \in \Gamma_{hol}(M, L^m) \mid \exists c \in \mathbb{C}^* : s = c \cdot e_\alpha^{(m)}\}$$

is a well-defined element of the projective space $\mathbb{P}(\Gamma_{hol}(M, L^m))$, only depending on $x = \tau(\alpha) \in M$.

This kind of coherent states go back to Berezin. A coordinate independent version and extensions to line bundles were given by Rawnsley [71]. It plays an important role in the work of Cahen, Gutt, and Rawnsley on the quantization of Kähler manifolds [26, 27, 28, 29], via Berezin's covariant symbols. I will return to this in Section 6.5. In these works the coherent

vectors are parameterized by the elements of $L \setminus 0$. The definition here uses the points of the total space of the dual bundle U . It has the advantage that one can consider all tensor powers of L together on an equal footing.

Definition 6.2. The *coherent state embedding* is the antiholomorphic embedding

$$M \rightarrow \mathbb{P}(\Gamma_{hol}(M, L^m)) \cong \mathbb{P}^N(\mathbb{C}), \quad x \mapsto [e_{\tau^{-1}(x)}^{(m)}]. \quad (6.4)$$

Here $N = \dim \Gamma_{hol}(M, L^m) - 1$. In this review, in abuse of notation, we will understand under $\tau^{-1}(x)$ always a non-zero element of the fiber over x . The coherent state embedding is up to conjugation the embedding of Section 2.9 with respect to an orthonormal basis of the sections. In [15] further results on the geometry of the coherent state embedding are given.

6.2. Covariant Berezin symbols. We start with the

Definition 6.3. The *covariant Berezin symbol* $\sigma^{(m)}(A)$ (of level m) of an operator $A \in \text{End}(\Gamma_{hol}(M, L^{(m)}))$ is defined as

$$\sigma^{(m)}(A) : M \rightarrow \mathbb{C}, \quad x \mapsto \sigma^{(m)}(A)(x) := \frac{\langle e_\alpha^{(m)}, A e_\alpha^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle}, \quad \alpha \in \tau^{-1}(x). \quad (6.5)$$

As the factors appearing in (6.3) will cancel, it is a well-defined function on M . If the level m is clear from the context I will sometimes drop it in the notation.

We consider also the *coherent projectors* used by Rawnsley

$$P_x^{(m)} = \frac{|e_\alpha^{(m)}\rangle \langle e_\alpha^{(m)}|}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle}, \quad \alpha \in \tau^{-1}(x). \quad (6.6)$$

Here we used the convenient bra-ket notation of the physicists. Recall, if s is a section then

$$P_x^{(m)} s = \frac{|e_\alpha^{(m)}\rangle \langle e_\alpha^{(m)}, s \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} = \frac{\langle e_\alpha^{(m)}, s \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} e_\alpha^{(m)}.$$

Again the projector is well-defined on M . With its help the covariant symbol can be expressed as

$$\sigma^{(m)}(A) = \text{Tr}(A P_x^{(m)}). \quad (6.7)$$

From the definition of the symbol it follows that $\sigma^{(m)}(A)$ is real analytic and

$$\sigma^{(m)}(A^*) = \overline{\sigma^{(m)}(A)}. \quad (6.8)$$

6.3. Rawnsley's ϵ function. Rawnsley [71] introduced a very helpful function on the manifold M relating the local metric in the bundle with the scalar product on coherent states. In our dual description we define it in the following way.

Definition 6.4. *Rawnsley's epsilon function* is the function

$$M \rightarrow C^\infty(M), \quad x \mapsto \epsilon^{(m)}(x) := \frac{h^{(m)}(e_\alpha^{(m)}, e_\alpha^{(m)})(x)}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle}, \quad \alpha \in \tau^{-1}(x). \quad (6.9)$$

With (6.3) it is clear that it is a well-defined function on M . Furthermore, using (6.1)

$$0 \neq \langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle = \alpha^{\otimes m}(e_\alpha^{(m)}(\tau(\alpha)))$$

it follows that

$$e_\alpha^{(m)}(x) \neq 0, \quad \text{for } x = \tau(\alpha), \quad \text{and } \epsilon^{(m)} > 0. \quad (6.10)$$

Hence, we can define the modified measure

$$\Omega_\epsilon^{(m)}(x) := \epsilon^{(m)}(x)\Omega(x) \quad (6.11)$$

for the space of functions on M and obtain a modified scalar product $\langle \cdot, \cdot \rangle_\epsilon^{(m)}$ for $C^\infty(M)$.

Proposition 6.5. *For $s_1, s_2 \in \Gamma_{hol}(M, L^m)$ we have*

$$\begin{aligned} h^{(m)}(s_1, s_2)(x) &= \frac{\overline{\langle e_\alpha^{(m)}, s_1 \rangle} \langle e_\alpha^{(m)}, s_2 \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \cdot \epsilon^{(m)}(x) \\ &= \langle s_1, P_x^{(m)} s_2 \rangle \cdot \epsilon^{(m)}(x). \end{aligned} \quad (6.12)$$

Proof. Due to (6.10) we can represent every section s locally at x as $s(x) = \hat{s}(x)e_\alpha^{(m)}$ with a local function $\hat{s}$. Now

$$\langle e_\alpha^{(m)}, s \rangle = \alpha^{(m)}(\hat{s}(x)e_\alpha^{(m)}(x)) = \hat{s}(x)\alpha^{(m)}(e_\alpha^{(m)}(x)) = \hat{s}(x)\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle.$$

We rewrite $h^{(m)}(s_1, s_2)(x) = \overline{\hat{s}_1} s_2 h^{(m)}(e_\alpha^{(m)}, e_\alpha^{(m)})(x)$, and obtain

$$h^{(m)}(s_1, s_2)(x) = \frac{\overline{\langle e_\alpha^{(m)}, s_1 \rangle}}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \frac{\langle e_\alpha^{(m)}, s_2 \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \cdot h^{(m)}(e_\alpha^{(m)}, e_\alpha^{(m)})(x). \quad (6.13)$$

From the definition (6.9) the first relation follows. Obviously, it can be rewritten with the coherent projector to obtain the second relation. $\square$

There exists another useful description of the epsilon function.

Proposition 6.6. *Let $s_1, s_2, \dots, s_k$ be an arbitrary orthonormal basis of $\Gamma_{hol}(M, L^m)$. Then*

$$\epsilon^{(m)}(x) = \sum_{j=1}^k h^{(m)}(s_j, s_j)(x). \quad (6.14)$$

Proof. For every vector ψ in a finite-dimensional hermitian vector space with orthonormal basis $s_j, j = 1, \dots, k$ the coefficient with respect to the basis element s_j is given by $\psi_j = \langle s_j, \psi \rangle$. Furthermore, $\langle \psi, \psi \rangle = \|\psi\|^2 = \sum_j \overline{\psi_j} \psi_j$. Now from (6.12)

$$\sum_{j=1}^k h(s_j, s_j)(x) = \frac{\epsilon^{(m)}(x)}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \sum_{j=1}^k \overline{\langle e_\alpha^{(m)}, s_j \rangle} \langle e_\alpha^{(m)}, s_j \rangle.$$

Hence the claim. $\square$

In certain special cases the functions $\epsilon^{(m)}$ will be constant as function of the points of the manifold. In this case we can apply Proposition 6.11 below for $A = id$, the identity operator, and obtain

$$\epsilon^{(m)} = \frac{\dim \Gamma_{hol}(M, L^m)}{\text{vol } M}. \quad (6.15)$$

Here $\text{vol } M$ denotes the volume of the manifold with respect to the Liouville measure. Now the question arises when $\epsilon^{(m)}$ will be constant, resp. when the measure $\Omega_\epsilon^{(m)}$ will be the standard measure (up to a scalar). From Proposition 6.6 it follows that if there is a transitive group action on the manifold and everything, e.g. Kähler form, bundle, metric, is homogenous with respect to the action this will be the case. An example is given by $M = \mathbb{P}^N(\mathbb{C})$. By a result of Rawnsley [71], resp. Cahen, Gutt and Rawnsley [26], $\epsilon^{(m)} \equiv \text{const}$ if and only if the quantization is projectively induced. This means that under the conjugate of the coherent state embedding, the Kähler form ω of M coincides with the pull-back of the Fubini-Study form. Note that in general this is not the case, see Section 7.6.

6.4. Contravariant Berezin symbols. Recall the modified Liouville measure (6.11) and modified scalar product for the functions on M introduced in the last subsection.

Definition 6.7. Given an operator $A \in \text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ then a *contravariant Berezin symbol* $\check{\sigma}^{(m)}(A) \in C^\infty(M)$ of A is defined by the representation of the operator A as integral

$$A = \int_M \check{\sigma}^{(m)}(A)(x) P_x^{(m)} \Omega_\epsilon^{(m)}(x), \quad (6.16)$$

if such a representation exists.

Proposition 6.8. The Toeplitz operator $T_f^{(m)}$ admits a representation (6.16) with

$$\check{\sigma}^{(m)}(T_f^{(m)}) = f, \quad (6.17)$$

i.e. the function f is a contravariant symbol of the Toeplitz operator $T_f^{(m)}$. Moreover, every operator $A \in \text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ has a contravariant symbol.

Proof. Let $f \in C^\infty(M)$ and set

$$A := \int_M f(x) P_x^{(m)} \Omega_\epsilon^{(m)}(x), \quad (6.18)$$

then $\check{\sigma}^{(m)}(A) = f$. For arbitrary $s_1, s_2 \in \Gamma_{\text{hol}}(M, L^m)$ we calculate (using (6.12))

$$\begin{aligned} \langle s_1, A s_2 \rangle &= \int_M f(x) \langle s_1, P_x^{(m)} s_2 \rangle \Omega_\epsilon^{(m)}(x) = \int_M f(x) h^{(m)}(s_1, s_2)(x) \Omega(x) \\ &= \int_M h^{(m)}(s_1, f s_2)(x) \Omega(x) = \langle s_1, f s_2 \rangle = \langle s_1, T_f^{(m)} s_2 \rangle. \end{aligned} \quad (6.19)$$

Hence $T_f^{(m)} = A$. As the Toeplitz map is surjective (Proposition 3.6) every operator is a Toeplitz operator, hence has a contravariant symbol. $\square$

Note that given an operator its contravariant symbol on a fixed level m is not uniquely defined.

We introduce on $\text{End}(\Gamma_{\text{hol}}(M, L^{(m)}))$ the Hilbert-Schmidt norm

$$\langle A, C \rangle_{HS} = \text{Tr}(A^* \cdot C). \quad (6.20)$$

Theorem 6.9. *The Toeplitz map $f \rightarrow T_f^{(m)}$ and the covariant symbol map $A \rightarrow \sigma^{(m)}(A)$ are adjoint:*

$$\langle A, T_f^{(m)} \rangle_{HS} = \langle \sigma^{(m)}(A), f \rangle_\epsilon^{(m)}. \quad (6.21)$$

Proof.

$$\langle A, T_f^{(m)} \rangle = \text{Tr}(A^* \cdot T_f^{(m)}) = \text{Tr}(A^* \int_M f(x) P_x^{(m)} \Omega_\epsilon^{(m)}(x)) = \int_M f(x) \text{Tr}(A^* \cdot P_x^{(m)}) \Omega_\epsilon^{(m)}(x). \quad (6.22)$$

Now applying the definition (6.7) and equation (6.8)

$$\langle A, T_f^{(m)} \rangle = \int_M f(x) \sigma^{(m)}(A^*) \Omega_\epsilon^{(m)}(x) = \int_M \overline{\sigma^{(m)}(A)}(x) f(x) \Omega_\epsilon^{(m)}(x) = \langle \sigma^{(m)}(A), f(x) \rangle_\epsilon^{(m)}. \quad (6.23)$$

□

As every operator has a contravariant symbol we can also conclude

$$\langle A, B \rangle_{HS} = \langle \sigma^{(m)}(A), \check{\sigma}^{(m)}(B) \rangle_\epsilon^{(m)}. \quad (6.24)$$

From Theorem 6.9 by using the surjectivity of the Toeplitz map we get

Proposition 6.10. *The covariant symbol map $\sigma^{(m)}$ is injective.*

Another application is the following

Proposition 6.11.

$$\text{Tr } A = \int_M \sigma^{(m)}(A) \Omega_\epsilon^{(m)}. \quad (6.25)$$

Proof. We use $Id = T_1$ and by (6.21) $\text{Tr } A = \langle A, Id \rangle_{HS} = \langle \sigma^{(m)}(A), 1 \rangle_\epsilon^{(m)}$. □

6.5. Berezin star product. Under certain very restrictive conditions Berezin covariant symbols can be used to construct a star product, called the *Berezin star product*. Recall that Proposition 6.10 says that the linear symbol map

$$\sigma^{(m)} : \text{End}(\Gamma_{hol}(M, L^{(m)})) \rightarrow C^\infty(M) \quad (6.26)$$

is injective. Its image is a subspace $\mathcal{A}^{(m)}$ of $C^\infty(M)$, called the subspace of covariant symbols of level m . If $\sigma^{(m)}(A)$ and $\sigma^{(m)}(B)$ are elements of this subspace the operators A and B will be uniquely fixed. Hence also $\sigma^{(m)}(A \cdot B)$. Now one takes

$$\sigma^{(m)}(A) \star_{(m)} \sigma^{(m)}(B) := \sigma^{(m)}(A \cdot B) \quad (6.27)$$

as definition for an associative and noncommutative product $\star_{(m)}$ on $\mathcal{A}^{(m)}$.

It is even possible to give an analytic expression for the resulting symbol. For this we introduce the *two-point function*

$$\psi^{(m)}(x, y) = \frac{\langle e_\alpha^{(m)}, e_\beta^{(m)} \rangle \langle e_\beta^{(m)}, e_\alpha^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle \langle e_\beta^{(m)}, e_\beta^{(m)} \rangle} \quad (6.28)$$

with $\alpha = \tau^{-1}(x) = x$ and $\beta = \tau^{-1}(y)$. This function is well-defined on $M \times M$. Furthermore, we have the *two-point symbol*

$$\sigma^{(m)}(A)(x, y) = \frac{\langle e_\alpha^{(m)}, A e_\beta^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\beta^{(m)} \rangle}. \quad (6.29)$$

It is the analytic extension of the real-analytic covariant symbol. It is well-defined on an open dense subset of $M \times M$ containing the diagonal. Using (6.12) we express

$$\begin{aligned} \sigma^{(m)}(A \cdot B)(x) &= \frac{\langle e_\alpha^{(m)}, A \cdot B e_\alpha^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} = \frac{\langle A^* e_\alpha^{(m)}, B e_\alpha^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \\ &= \int_M h^{(m)}(A^* e_\alpha^{(m)}, B e_\alpha^{(m)})(y) \frac{\Omega(y)}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} = \int_M \frac{\langle e_\alpha^{(m)}, A e_\beta^{(m)} \rangle \langle e_\beta^{(m)}, B e_\alpha^{(m)} \rangle}{\langle e_\beta^{(m)}, e_\beta^{(m)} \rangle} \frac{\epsilon^{(m)}(y) \Omega(y)}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \\ &= \int_M \sigma^{(m)}(A)(x, y) \sigma^{(m)}(B)(y, x) \cdot \psi^{(m)}(x, y) \cdot \epsilon^{(m)}(y) \Omega(y). \quad (6.30) \end{aligned}$$

The crucial problem is how to relate different levels m to define for all possible symbols a unique product not depending on m . In certain special situations like these studied by Berezin himself [17] and Cahen, Gutt, and Rawnsley [26] the subspaces are nested into each other and the union $\mathcal{A} = \bigcup_{m \in \mathbb{N}} \mathcal{A}^{(m)}$ is a dense subalgebra of $C^\infty(M)$. Indeed, in the cases considered, the manifold is a homogenous manifold and the epsilon function $\epsilon^{(m)}$ is a constant. A detailed analysis shows that in this case a star product is given.

For further examples, for which this method works (not necessarily compact) see other articles by Cahen, Gutt, and Rawnsley [27, 28, 29]. For related results see also work of Moreno and Ortega-Navarro [67], [66]. In particular, also the work of Engliš [42, 41, 40, 39]. Reshetikhin and Takhtajan [72] gave a construction of a (formal) star product using formal integrals in the spirit of the Berezin's covariant symbol construction.

7. BEREZIN TRANSFORM

7.1. The definition. Starting from $f \in C^\infty(M)$ we can assign to it its Toeplitz operator $T_f^{(m)} \in \text{End}(\Gamma_{hol}(M, L^{(m)}))$ and then assign to $T_f^{(m)}$ the covariant symbol $\sigma^{(m)}(T_f^{(m)})$. It is again an element of $C^\infty(M)$.

Definition 7.1. The map

$$C^\infty(M) \rightarrow C^\infty(M), \quad f \mapsto I^{(m)}(f) := \sigma^{(m)}(T_f^{(m)}) \quad (7.1)$$

is called the *Berezin transform (of level m)*.

From the point of view of Berezin's approach the operator $T_f^{(m)}$ has as a contravariant symbol f . Hence $I^{(m)}$ gives a correspondence between contravariant symbols and covariant symbols of operators. The Berezin transform was introduced and studied by Berezin [17] for certain classical symmetric domains in $\mathbb{C}^n$. These results were extended by Unterberger and Upmeyer [85], see also Engliš [40, 41, 42] and Engliš and Peetre [43]. Obviously, the Berezin transform makes perfect sense in the compact Kähler case which we consider here.

7.2. The asymptotic expansion. The results presented here are joint work with Alexander Karabegov [58]. See also [79] for an overview.

Theorem 7.2. *Given $x \in M$ then the Berezin transform $I^{(m)}(f)$ evaluated at the point x has a complete asymptotic expansion in powers of $1/m$ as $m \rightarrow \infty$*

$$I^{(m)}(f)(x) \sim \sum_{i=0}^{\infty} I_i(f)(x) \frac{1}{m^i}, \quad (7.2)$$

where $I_i : C^\infty(M) \rightarrow C^\infty(M)$ are maps with

$$I_0(f) = f, \quad I_1(f) = \Delta f. \quad (7.3)$$

Here the Δ is the usual Laplacian with respect to the metric given by the Kähler form ω .

Complete asymptotic expansion means the following. Given $f \in C^\infty(M)$, $x \in M$ and an $r \in \mathbb{N}$ then there exists a positive constant A such that

$$\left| I^{(m)}(f)(x) - \sum_{i=0}^{r-1} I_i(f)(x) \frac{1}{m^i} \right|_\infty \leq \frac{A}{m^r}. \quad (7.4)$$

In Section 7.4 I will give some remarks on the proof. But before I present you a nice application

7.3. Norm preservation of the BT operators. In [77] I conjectured (7.2) (which is now a mathematical result) and showed how such an asymptotic expansion supplies a different proof of Theorem 3.3, part (a). For completeness I reproduce the proof here.

Proposition 7.3.

$$|I^{(m)}(f)|_\infty = |\sigma^{(m)}(T_f^{(m)})|_\infty \leq \|T_f^{(m)}\| \leq |f|_\infty. \quad (7.5)$$

Proof. Using Cauchy-Schwarz inequality we calculate ($x = \tau(\alpha)$)

$$|\sigma^{(m)}(T_f^{(m)})(x)|^2 = \frac{|\langle e_\alpha^{(m)}, T_f^{(m)} e_\alpha^{(m)} \rangle|^2}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle^2} \leq \frac{\langle T_f^{(m)} e_\alpha^{(m)}, T_f^{(m)} e_\alpha^{(m)} \rangle}{\langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle} \leq \|T_f^{(m)}\|^2. \quad (7.6)$$

Here the last inequality follows from the definition of the operator norm. This shows the first inequality in (7.5). For the second inequality introduce the multiplication operator $M_f^{(m)}$ on $\Gamma_\infty(M, L^m)$. Then $\|T_f^{(m)}\| = \|\Pi^{(m)} M_f^{(m)} \Pi^{(m)}\| \leq \|M_f^{(m)}\|$ and for $\varphi \in \Gamma_\infty(M, L^m)$, $\varphi \neq 0$

$$\frac{\|M_f^{(m)} \varphi\|^2}{\|\varphi\|^2} = \frac{\int_M h^{(m)}(f\varphi, f\varphi) \Omega}{\int_M h^{(m)}(\varphi, \varphi) \Omega} = \frac{\int_M f(z) \overline{f(z)} h^{(m)}(\varphi, \varphi) \Omega}{\int_M h^{(m)}(\varphi, \varphi) \Omega} \leq |f|_\infty^2. \quad (7.7)$$

Hence,

$$\|T_f^{(m)}\| \leq \|M_f^{(m)}\| = \sup_{\varphi \neq 0} \frac{\|M_f^{(m)} \varphi\|}{\|\varphi\|} \leq |f|_\infty. \quad (7.8)$$

□

Proof. (Theorem 3.3 part (a).) Choose as $x_e \in M$ a point with $|f(x_e)| = |f|_\infty$. From the fact that the Berezin transform has as leading term the identity it follows that $|(I^{(m)}f)(x_e) - f(x_e)| \leq C/m$ with a suitable constant C . Hence, $||f(x_e)| - |(I^{(m)}f)(x_e)|| \leq C/m$ and

$$|f|_\infty - \frac{C}{m} = |f(x_e)| - \frac{C}{m} \leq |(I^{(m)}f)(x_e)| \leq |I^{(m)}f|_\infty. \quad (7.9)$$

Putting (7.5) and (7.9) together we obtain

$$|f|_\infty - \frac{C}{m} \leq \|T_f^{(m)}\| \leq |f|_\infty. \quad (7.10)$$

□

7.4. Bergman kernel. To understand the Berezin transform better we have to study the Bergman kernel. Recall from Section 5 the Szegő projectors $\Pi : L^2(Q, \mu) \rightarrow \mathcal{H}$ and its components $\hat{\Pi}^{(m)} : L^2(Q, \mu) \rightarrow \mathcal{H}^{(m)}$, the Bergman projectors. The Bergman projectors have smooth integral kernels, the *Bergman kernels* $\mathcal{B}_m(\alpha, \beta)$ defined on $Q \times Q$, i.e.

$$\hat{\Pi}^{(m)}(\psi)(\alpha) = \int_Q \mathcal{B}_m(\alpha, \beta) \psi(\beta) \mu(\beta). \quad (7.11)$$

The Bergman kernels can be expressed with the help of the coherent vectors.

Proposition 7.4.

$$\mathcal{B}_m(\alpha, \beta) = \psi_{e_\beta^{(m)}}(\alpha) = \overline{\psi_{e_\alpha^{(m)}}(\beta)} = \langle e_\alpha^{(m)}, e_\beta^{(m)} \rangle. \quad (7.12)$$

For the proofs of this and the following propositions see [58], or [79].

Let $x, y \in M$ and choose $\alpha, \beta \in Q$ with $\tau(\alpha) = x$ and $\tau(\beta) = y$ then the functions

$$u_m(x) := \mathcal{B}_m(\alpha, \alpha) = \langle e_\alpha^{(m)}, e_\alpha^{(m)} \rangle, \quad (7.13)$$

$$v_m(x, y) := \mathcal{B}_m(\alpha, \beta) \cdot \mathcal{B}_m(\beta, \alpha) = \langle e_\alpha^{(m)}, e_\beta^{(m)} \rangle \cdot \langle e_\beta^{(m)}, e_\alpha^{(m)} \rangle \quad (7.14)$$

are well-defined on M and on $M \times M$ respectively. The following proposition gives an integral representation of the Berezin transform.

Proposition 7.5.

$$\begin{aligned} \left(I^{(m)}(f) \right) (x) &= \frac{1}{\mathcal{B}_m(\alpha, \alpha)} \int_Q \mathcal{B}_m(\alpha, \beta) \mathcal{B}_m(\beta, \alpha) \tau^* f(\beta) \mu(\beta) \\ &= \frac{1}{u_m(x)} \int_M v_m(x, y) f(y) \Omega(y). \end{aligned} \quad (7.15)$$

Typically, asymptotic expansions can be obtained using stationary phase integrals. But for such an asymptotic expansion of the integral representation of the Berezin transform we will not only need an asymptotic expansion of the Bergman kernel along the diagonal (which is well-known) but in a neighbourhood of it. This is one of the key results obtained in [58]. It is based on works of Boutet de Monvel and Sjöstrand [25] on the Szegő kernel and in generalization of a result of Zelditch [92] on the Bergman kernel on the diagonal. The integral representation is used then to prove the existence of the asymptotic expansion of the Berezin transform.

Having such an asymptotic expansion it still remains to identify its terms. As it was explained in Section 4.8, Karabegov assigns to every formal deformation quantization with the “separation of variables” property a *formal Berezin transform* I . In [58] it is shown that there is an explicitly specified star product $\star$ (see Theorem 5.9 in [58]) with associated formal Berezin transform such that if we replace $\frac{1}{m}$ by the formal variable ν in the asymptotic expansion of the Berezin transform $I^{(m)}f(x)$ we obtain $I(f)(x)$. This finally proves Theorem 7.2. We will exhibit the star product $\star$ in the next section.

7.5. Identification of the BT star product. Moreover in [58] there is another object introduced, the *twisted product*

$$R^{(m)}(f, g) := \sigma^{(m)}(T_f^{(m)} \cdot T_g^{(m)}) . \quad (7.16)$$

Also for it the existence of a complete asymptotic expansion was shown. It was identified with a twisted formal product. This allows the identification of the BT star product with a special star product within the classification of Karabegov. From this identification the properties of Theorem 4.6 of locality, separation of variables type, and the calculation to the classifying forms and classes for the BT star product follows.

As already announced in Section 4.8, the BT star product $\star_{BT}$ is the opposite of the dual star product of a certain star product $\star$. To identify $\star$ we will give its classifying Karabegov form $\widehat{\omega}$. As already mentioned above, Zelditch [92] proved that the function u_m (7.13) has a complete asymptotic expansion in powers of $1/m$. In detail he showed

$$u_m(x) \sim m^n \sum_{k=0}^{\infty} \frac{1}{m^k} b_k(x), \quad b_0 = 1. \quad (7.17)$$

If we replace in the expansion $1/m$ by the formal variable ν we obtain a formal function s defined by

$$e^s(x) = \sum_{k=0}^{\infty} \nu^k b_k(x). \quad (7.18)$$

Now take as formal potential (4.16)

$$\widehat{\Phi} = \frac{1}{\nu} \Phi_{-1} + s,$$

where Φ_{-1} is the local Kähler potential of the Kähler form $\omega = \omega_{-1}$. Then $\widehat{\omega} = i\partial\bar{\partial}\widehat{\Phi}$. It might be also written in the form

$$\widehat{\omega} = \frac{1}{\nu} \omega + \mathbb{F}(i\partial\bar{\partial} \log \mathcal{B}_m(\alpha, \alpha)). \quad (7.19)$$

Here we denote the replacement of $1/m$ by the formal variable ν by the symbol $\mathbb{F}$.

7.6. Pullback of the Fubini-Study form. Starting from the Kähler manifold (M, ω) and after choosing an orthonormal basis of the space $\Gamma_{hol}(M, L^m)$ we obtain an embedding

$$\phi^{(m)} : M \rightarrow \mathbb{P}^{N(m)}$$

of M into projective space of dimension $N(m)$. On $\mathbb{P}^{N(m)}$ we have the standard Kähler form, the Fubini-Study form ω_{FS} . The pull-back $(\phi^{(m)})^* \omega_{FS}$ will not depend on the orthogonal

basis chosen for the embedding. But in general it will not coincide with a scalar multiple of the Kähler form ω we started with (see [15] for a thorough discussion of the situation).

It was shown by Zelditch [92], by generalizing a result of Tian [83], that $(\Phi^{(m)})^* \omega_{FS}$ admits a complete asymptotic expansion in powers of $\frac{1}{m}$ as $m \rightarrow \infty$. In fact it is related to the asymptotic expansion of the Bergman kernel (7.13) along the diagonal. The pull-back can be given as [92, Prop.9]

$$\left(\phi^{(m)}\right)^* \omega_{FS} = m\omega + i\partial\bar{\partial} \log u_m(x). \quad (7.20)$$

If we again replace $1/m$ by ν we obtain via (7.19) the Karabegov form introduced in Section 4.8

$$\hat{\omega} = \mathbb{F}\left(\left(\phi^{(m)}\right)^* \omega_{FS}\right). \quad (7.21)$$

REFERENCES

- [1] Abraham, R., and Marsden, J.E., *Foundations of Mechanics*. Addison Wesley, Redwood City, 1985.
- [2] Ali, S.T., *Quantization techniques: A quick overview*. in Contemporary Problems in Mathematical Physics, World Scientific 2002.
- [3] Ali, S.T., Doebner, H.D., *Ordering problem in quantum mechanics: Prime quantization and a physical interpretation*. Phys. Rev. **A 41** (1990), 1199–1210.
- [4] Ali, S.T., Engliš, M., *Quantization methods: A guide for physicists and analysts*. Rev. Math. Phys. **17**(4) (2005), 391–490.
- [5] Andersen, J.E., *Asymptotic faithfulness of the quantum $SU(n)$ representations of the mapping class group*. Ann. of Math. **163** (2006), 347–368.
- [6] Andersen, J.E., *Deformation quantization and geometric quantization of abelian moduli spaces*. Comm. Math. Phys. **255** (2005), 727–745.
- [7] Andersen, J.E., *Mapping class groups do not have property T*. arXiv:0706.2184.
- [8] Andersen, J.E., and Gammelgaard, N.L., *Hitchin’s projectively flat connection, Toeplitz operators and the asymptotic expansion of TQFT curve operators*. arXiv:0903.4091.
- [9] Andersen, J.E., Ueno, K., *Abelian Conformal field theories and the determinant bundles*. Internat. Jour. of Math. **18** (2007), 919–993.
- [10] Andersen, J.E., Ueno, K., *Constructing modular functors from conformal field theories*. Jour. of Knot Theories and its Ramifications **16** (2007), 127–202.
- [11] Aschieri, P., Madore, J., Manousselis, Zoupanes, G., *Dimensional reduction over fuzzy coset spaces*. JHEP **04** (2004), 034.
- [12] Balachandran, A.P., Dolan, B.P., Lee, J.H., Martin, X., O’Connor, D., *Fuzzy complex projective spaces and their star-products*. JGP **43** (2002), 184.
- [13] Balachandran, A.P., Kurkuoglu, S., Vaidya, S., *Lectures on fuzzy and fuzzy SUSY physics*. hep-th/0511114.
- [14] Bayen, F., Flato, M., Fronsdal, C., Lichnerowicz, A., and Sternheimer, D., *Deformation theory and quantization, Part I*. Lett. Math. Phys. **1** (1977), 521–530: *Deformation theory and quantization, Part II and III*. Ann. Phys. **111** (1978), 61–110, 111–151.
- [15] Berceanu, St., and Schlichenmaier, M., *Coherent state embeddings, polar divisors and Cauchy formulas*. JGP **34** (2000), 336–358.
- [16] Berezin, F.A., *Quantization*. Math. USSR-Izv. **8** (1974), 1109–1165.
- [17] Berezin, F.A., *Quantization in complex symmetric spaces*. Math. USSR-Izv. **9** (1975), 341–379.
- [18] Berger, C.A., and Coburn, L.A., *Toeplitz operators and quantum mechanics*. J. Funct. Anal. **68** (1986), 273–299.
- [19] Bertelson, M., Cahen, M., and Gutt, S., *Equivalence of star products*. Classical Quantum Gravity **14** (1997), A93–A107.
- [20] Bordemann, M., Hoppe, J., Schaller, P., and Schlichenmaier, M., *$gl(\infty)$ and geometric quantization*. Commun. Math. Phys. **138** (1991), 209–244.

- [21] Bordemann, M., Meinrenken, E., and Schlichenmaier, M., *Toeplitz quantization of Kähler manifolds and $gl(n)$, $n \rightarrow \infty$ limits*. Commun. Math. Phys. **165** (1994), 281–296.
- [22] Bordemann, M., and Waldmann, St., *A Fedosov star product of the Wick type for Kähler manifolds*. Lett. Math. Phys. **41** (1997), 243–253.
- [23] Borthwick, D., Paul, D., and Uribe, A.: *Semiclassical spectral estimates for Toeplitz operators*. Ann. Inst. Fourier, Grenoble **48** (1998), 1189–1229.
- [24] Boutet de Monvel, L., and Guillemin, V., *The spectral theory of Toeplitz operators*. Ann. Math. Studies, Nr.99, Princeton University Press, Princeton, 1981.
- [25] Boutet de Monvel, L., and Sjostrand, J., *Sur la singularité des noyaux de Bergman et de Szegő*. Asterisque **34-35** (1976), 123–164.
- [26] Cahen, M., Gutt, S., and Rawnsley, J., *Quantization of Kähler manifolds I: Geometric interpretation of Berezin’s quantization*. JGP **7** (1990), 45–62.
- [27] Cahen, M., Gutt, S., and Rawnsley, J., *Quantization of Kähler manifolds II*. Trans. Amer. Math. Soc. **337** (1993), 73–98.
- [28] Cahen, M., Gutt, S., and Rawnsley, J., *Quantization of Kähler manifolds III*. Lett. Math. Phys. **30** (1994), 291–305.
- [29] Cahen, M., Gutt, S., and Rawnsley, J., *Quantization of Kähler manifolds IV*. Lett. Math. Phys. **34** (1995), 159–168.
- [30] Charles, L., *Berezin-Toeplitz operators, a semi-classical approach*. Comm. Math. Phys. **239** (2003), 1–28.
- [31] Charles, L., *Toeplitz operators and Hamiltonian torus actions*. Jour. Functional Anal. **236** (2006), 299–350.
- [32] Charles, L., *Semi-classical properties of geometric quantization with metaplectic corrections*. Comm. Math. Phys. **270** (2007), 445–480.
- [33] Carow-Watamura, U., Steinacker, H., Watamura, S., *Monopole bundles over fuzzy complex projective spaces*. JGP **54** (2005), 373.
- [34] Coburn, L.A., *Deformation estimates for the Berezin-Toeplitz quantization*. Commun. Math. Phys. **149** (1992), 415–424.
- [35] De Wilde, M., and Lecomte, P.B.A., *Existence of star products and of formal deformations of the Poisson-Lie algebra of arbitrary symplectic manifolds*. Lett. Math. Phys. **7** (1983), 487–496.
- [36] Deligne, P., *Deformation de l’algebre des fonctions d’une variete symplectique: Comparaison entre Fedosov et De Wilde, Lecomte*. Sel. Math., New Ser. **1** (1995), 667–697.
- [37] Dito, G., and Sternheimer, D., *Deformation quantization: genesis, developments, and metamorphoses*. (in) IRMA Lectures in Math. Theoret. Phys. Vol 1, Walter de Gruyter, Berlin 2002, 9–54, math.QA/0201168.
- [38] Dolan, B.P., Huet, I., Murray, S., O’Connor, D., *Noncommutative vector bundles over fuzzy $\mathbb{CP}^N$ and their covariant derivatives*. JHEP **07** (2007), 007.
- [39] Engliš, M., *The asymptotics of a Laplace integral on a Kähler manifold*. J. Reine Angew. Math. **528** (2000), 1–39.
- [40] Engliš, M., *Asymptotics of the Berezin transform and quantization on planar domains*. Duke Math. J. **79** (1995), 57–76.
- [41] Engliš, M., *Berezin quantization and reproducing kernels on complex domains*. Trans. Amer. Math. Soc. **348** (1996), 411–479.
- [42] Engliš, M., *Weighted Bergman kernels and quantization*. Commun. Math. Phys. **227** (2002), 211–241.
- [43] Engliš, M., and Peetre, J., *On the correspondence principle for the quantized annulus*. Math. Scand. **78** (1996), 183–206.
- [44] Fedosov, B.V., *Deformation quantization and index theory*. Akademie Verlag, Berlin, 1996.
- [45] Fedosov, B.V., *Deformation quantization and asymptotic operator representation*, Funktional Anal. i. Prilozhen. **25** (1990), 184–194; *A simple geometric construction of deformation quantization*. J. Diff. Geo. **40** (1994), 213–238.
- [46] Grosse, H., Maceda, M., Madore, J., Steinacker, H., *Fuzzy instantons*. Int. J. Mod. Phys. **A 17** (2002), 2095.
- [47] Grosse, H., Steinacker, H., *Finite gauge theory on fuzzy $\mathbb{CP}^2$* . Nucl. Phys. **B 704** (2005), 145.
- [48] Guillemin, V., *Some classical theorems in spectral theory revisited*. Seminars on singularities of solutions of linear partial differential equations, Ann. Math. Studies, Nr.91 (L. Hörmander, ed.), Princeton University Press, 1979, pp. 219–259.

- [49] Guillemin, V., *Star products on pre-quantizable symplectic manifolds*, Lett. Math. Phys. **35** (1995), 85–89.
- [50] Gutt, S., Rawnsley, J., *Equivalence of star products on a symplectic manifold: an introduction to Deligne’s Čech cohomology classes*. JGP **29**, No.4, 347–392 (1999).
- [51] Hawkins, E., *The Correspondence between Geometric Quantization and Formal Deformation Quantization*. math/9811049
- [52] Hörmander, L., *The Analysis of Linear Partial Differential Operators I, Distribution Theory and Fourier Analysis*. Springer Verlag, New York, 1983.
- [53] Karabegov, A.V., *Deformation quantization with separation of variables on a Kähler manifold*, Commun. Math. Phys. **180** (1996), 745–755.
- [54] Karabegov, A.V., *Cohomological classification of deformation quantizations with separation of variables*. Lett. Math. Phys. **43** (1998), 347–357.
- [55] Karabegov, A.V., *On the canonical normalization of a trace density of deformation quantization*. Lett. Math. Phys. **45** (1998), 217–228.
- [56] Karabegov, A.V., *Pseudo-Kähler quantization on flag manifolds* Commun. Math. Phys. **200** (1999), 355–379.
- [57] Karabegov, A.V., *On Fedosov’s approach to deformation quantization with separation of variables*, (in) Proceedings of the Conference Moshe Flato 1999, (eds. G. Dito, and D. Sternheimer), Kluwer 2000, 167–176.
- [58] Karabegov, A.V., Schlichenmaier, M., *Identification of Berezin-Toeplitz deformation quantization*. J. reine angew. Math. **540** (2001), 49–76
- [59] Klimek, S., and Lesniewski, A., *Quantum Riemann surfaces: I. The unit disc*. Comm. Math. Phys. **146** (1992) 103–122; *Quantum Riemann surfaces: II. The discrete series*. Lett. Math. Phys. **24** (1992) 125–139.
- [60] Kontsevich, M., *Deformation quantization of Poisson manifolds*. Lett. Math. Phys. **66** (2003), 157–216, preprint q-alg/9709040.
- [61] Landsman, N.P., *Mathematical topics between classical and quantum mechanics*. Springer, New York, Berlin, Heidelberg, 1998.
- [62] Lazaroiu, C.I., McNamee, D., Sämann, *Generalized Berezin quantization, Bergman metric and fuzzy laplacians*. JHEP, Sept. 2008.
- [63] Ma, X., and Marinescu, G., *Holomorphic Morse inequalities and Bergman kernels*. Birkhäuser, Basel, Boston, Berlin, 2007.
- [64] Ma, X., and Marinescu, G., *Toeplitz operators on symplectic manifolds*. J. Geom. Anal. **18** (2008), 565–611.
- [65] Madore, J., *The fuzzy sphere*. Class. and Quant. Grav. **9** (1992), 69.
- [66] Moreno, C., **-products on some Kähler manifolds*, Lett. Math. Phys. **11** (1986), 361–372.
- [67] Moreno, C., and Ortega-Navarro, P., **-products on $D^1(\mathbb{C})$, S^2 and related spectral analysis*, Lett. Math. Phys. **7** (1983), 181–193.
- [68] Moyal, J., *Quantum mechanics as a statistical theory*. Proc. Camb. Phil. Soc. **45** (1949), 99–124.
- [69] Nest, R., and Tsygan, B., *Algebraic index theory*. Commun. Math. Phys. **172** (1995), 223–262; *Algebraic index theory for families*. Advances in Math **113** (1995), 151–205.
- [70] Omori, H., Maeda, Y., and Yoshioka, A., *Weyl manifolds and deformation quantization*. Advances in Math **85** (1991), 224–255; *Existence of closed star-products*. Lett. Math. Phys. **26** (1992), 284–294.
- [71] Rawnsley, J.H., *Coherent states and Kähler manifolds*. Quart. J. Math. Oxford Ser.(2) **28** (1977), 403–415.
- [72] Reshetikhin, N., and Takhtajan, L., *Deformation quantization of Kähler manifolds*. Amer. Math. Soc. Transl (2) **201** (2000), 257–276.
- [73] Rieffel, M., *Questions on quantization*, Operator algebras and operator theory. Shanghai, China, July 4–9, 1997 (Liming et al., ed.), AMS, 1998, pp. 315–326.
- [74] Schlichenmaier, M., *Zwei Anwendungen algebraisch-geometrischer Methoden in der theoretischen Physik: Berezin-Toeplitz-Quantisierung und globale Algebren der zweidimensionalen konformen Feldtheorie*. Habilitationsschrift Universität Mannheim, 1996.
- [75] Schlichenmaier, M., *Berezin-Toeplitz quantization of compact Kähler manifolds*. in: Quantization, Coherent States and Poisson Structures, Proc. XIVth Workshop on Geometric Methods in Physics (Białowieża, Poland, 9–15 July 1995) (A, Strasburger, S.T. Ali, J.-P. Antoine, J.-P. Gazeau, and A. Odziejewicz, eds.), Polish Scientific Publisher PWN, 1998, q-alg/9601016, pp. 101–115.

- [76] Schlichenmaier, M., *Deformation quantization of compact Kähler manifolds via Berezin-Toeplitz operators*. Proceedings of the XXI Int. Coll. on Group Theoretical Methods in Physics (15-20 July 1996, Goslar, Germany) (H.-D. Doebner, P. Nattermann, and W. Scherer, eds.), World Scientific, 1997, pp. 396–400.
- [77] Schlichenmaier, M., *Berezin-Toeplitz quantization and Berezin symbols for arbitrary compact Kähler manifolds*. in Proceedings of the XVIIth workshop on geometric methods in physics, Białowieża, Poland, July 3 - 10, 1998 (eds. Schlichenmaier, et. al), (math.QA/9902066), Warsaw University Press, 45-56.
- [78] Schlichenmaier, M., *Deformation quantization of compact Kähler manifolds by Berezin-Toeplitz quantization*, (in) the Proceedings of the Conference Moshe Flato 1999, (eds. G.Dito, and D. Sternheimer), Kluwer 2000, 289 – 306, math.QA/9910137.
- [79] Schlichenmaier, M., *Berezin-Toeplitz quantization and Berezin transform*. (in) Long time behaviour of classical and quantum systems. Proc. of the Bologna APTEX Int. Conf. 13-17 September 1999, eds. S. Graffi, A. Martinez, World-Scientific, 2001, 271-287.
- [80] Schlichenmaier, M., *Berezin-Toeplitz quantization of the moduli space of flat $SU(N)$ connections*. J. Geom. Symmetry Phys. **9** (2007), 33-44.
- [81] Schlichenmaier, M., *An Introduction to Riemann surfaces, algebraic curves and moduli spaces*. 2nd ed., Springer, Berlin, Heidelberg, 2007.
- [82] Sternheimer, D., *Deformation quantization. Twenty years after*. In Particles, Fields and Gravitation. Lodz, 1998, AIP Conf. Proc. Vol 453.
- [83] Tian, G., *On a set of polarized Kähler metrics on algebraic manifolds*. J. Differential Geom. **32** (1990), 99–130.
- [84] Tuynman, G.M., *Generalized Bergman kernels and geometric quantization*. J. Math. Phys. **28**, (1987) 573–583.
- [85] Unterberger, A., and Upmeyer, H., *The Berezin transform and invariant differential operators*. Commun. Math. Phys. **164** (1994), 563–597.
- [86] Upmeyer, H., *Toeplitz operators on symmetric Siegel domains*. Math. Ann. **271** (1985), 401–414.
- [87] Upmeyer, H., *Toeplitz C^* -algebras on bounded symmetric domains*. Ann. of Math. **119** (1984), 549–576.
- [88] Upmeyer, H., *Fredholm indices for Toeplitz operators on bounded symmetric domains*. Am. J. Math. **110** (1988), 811–832.
- [89] Upmeyer, H., *Toeplitz C^* -algebras and noncommutative duality*. J. Oper. Theory **26** (1991), 407–432.
- [90] Upmeyer, H., *Toeplitz operators and index theory in several complex variables*. Birkhäuser, Basel 1996.
- [91] Weyl, H., *Gruppentheorie und Quantenmechanik*. 1931, Leipzig, Nachdruck Wissenschaftliche Buchgesellschaft, Darmstadt, 1977.
- [92] Zelditch, S., *Szegő kernels and a theorem of Tian*. Int. Math. Res. Not. **6** (1998), 317–331.

(Martin Schlichenmaier) UNIVERSITY OF LUXEMBOURG, MATHEMATICS RESEARCH UNIT, FSTC, 6, RUE COUDENHOVE-KALERGI L-1359 LUXEMBOURG-KIRCHBERG, GRAND DUCHY OF LUXEMBOURG

E-mail address: martin.schlichenmaier@uni.lu